

TABLE 6
Long-Term Trends in Thirty-Day Prevalence of Use of Various Drugs for Twelfth Graders

	Percent who used in last thirty days																									'99-'00 change		
	Class of 1975	Class of 1976	Class of 1977	Class of 1978	Class of 1979	Class of 1980	Class of 1981	Class of 1982	Class of 1983	Class of 1984	Class of 1985	Class of 1986	Class of 1987	Class of 1988	Class of 1989	Class of 1990	Class of 1991	Class of 1992	Class of 1993	Class of 1994	Class of 1995	Class of 1996	Class of 1997	Class of 1998	Class of 1999		Class of 2000	
<i>Approx. N</i> <i>(in thousands) =</i>	9.4	15.4	17.1	17.8	15.5	15.9	17.5	17.7	16.3	15.9	16.0	15.2	16.3	16.3	16.7	15.2	15.0	15.8	16.3	15.4	15.4	14.3	15.4	15.2	13.6	12.8		
Any Illicit Drug ^{a,b}	30.7	34.2	37.6	38.9	38.9	37.2	36.9	32.5	30.5	29.2	29.7	27.1	24.7	21.3	19.7	17.2	16.4	14.4	18.3	21.9	23.8	24.6	26.2	25.6	25.9	24.9	-1.0	
Any Illicit Drug Other Than Marijuana ^{a,b}	15.4	13.9	15.2	15.1	16.8	18.4	21.7	17.0	15.4	15.1	14.9	13.2	11.6	10.0	9.1	8.0	7.1	6.3	7.9	8.8	10.0	9.5	10.7	10.7	10.4	10.4	0.0	
Marijuana/Hashish	27.1	32.2	35.4	37.1	36.5	33.7	31.6	28.5	27.0	25.2	25.7	23.4	21.0	18.0	16.7	14.0	13.8	11.9	15.5	19.0	21.2	21.9	23.7	22.8	23.1	21.6	-1.6	
Inhalants ^c	—	0.9	1.3	1.5	1.7	1.4	1.5	1.5	1.7	1.9	2.2	2.5	2.8	2.6	2.3	2.7	2.4	2.3	2.5	2.7	3.2	2.5	2.5	2.3	2.0	2.2	+0.2	
Inhalants, Adjusted ^d	—	—	—	—	3.2	2.7	2.5	2.5	2.5	2.6	3.0	3.2	3.5	3.0	2.7	2.9	2.6	2.5	2.8	2.9	3.5	2.9	2.9	3.1	2.4	2.4	0.0	
Amyl/Butyl Nitrites ^{e,f}	—	—	—	—	2.4	1.8	1.4	1.1	1.4	1.4	1.6	1.3	1.3	0.6	0.6	0.6	0.4	0.3	0.6	0.4	0.4	0.7	0.7	1.0	0.4	0.3	-0.1	
Hallucinogens	4.7	3.4	4.1	3.9	4.0	3.7	3.7	3.4	2.8	2.6	2.5	2.5	2.2	2.2	2.2	2.2	2.1	2.7	3.1	4.4	3.5	3.9	3.8	3.5	2.6	-0.9ss		
Hallucinogens, Adjusted ^g	—	—	—	—	5.3	4.4	4.5	4.1	3.5	3.2	3.8	3.5	2.8	2.3	2.9	2.3	2.4	2.3	3.3	3.2	4.6	3.8	4.1	4.1	3.9	3.0	-0.9s	
LSD	2.3	1.9	2.1	2.1	2.4	2.3	2.5	2.4	1.9	1.5	1.6	1.7	1.8	1.8	1.8	1.9	1.9	2.0	2.4	2.6	4.0	2.5	3.1	3.2	2.7	1.6	-1.2sss	
Hallucinogens Other Than LSD	3.7	2.3	3.0	2.7	2.4	2.3	2.1	1.7	1.5	1.6	1.3	1.3	1.1	0.7	0.8	0.8	0.7	0.5	0.8	1.2	1.3	1.6	1.7	1.6	1.6	1.7	+0.1	
PCP ^{e,f}	—	—	—	—	2.4	1.4	1.4	1.0	1.3	1.0	1.6	1.3	0.6	0.3	1.4	0.4	0.5	0.6	1.0	0.7	0.6	1.3	0.7	1.0	0.8	0.9	+0.1	
MDMA (Ecstasy) ^e	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2.0	1.6	1.5	2.5	3.6	+1.1	
Cocaine	1.9	2.0	2.9	3.9	5.7	5.2	5.8	5.0	4.9	5.8	6.7	6.2	4.3	3.4	2.8	1.9	1.4	1.3	1.3	1.5	1.8	2.0	2.3	2.4	2.6	2.1	-0.5	
Crack ^h	—	—	—	—	—	—	—	—	—	—	—	—	1.3	1.6	1.4	0.7	0.7	0.6	0.7	0.8	1.0	1.0	0.9	1.0	1.1	1.0	-0.1	
Other Cocaine ⁱ	—	—	—	—	—	—	—	—	—	—	—	—	4.1	3.2	1.9	1.7	1.2	1.0	1.2	1.3	1.3	1.6	2.0	2.0	2.5	1.7	-0.7s	
Heroin ^j	0.4	0.2	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.2	0.2	0.2	0.3	0.2	0.2	0.3	0.2	0.3	0.6	0.5	0.5	0.5	0.5	0.7	+0.2	
With a needle ^k	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.3	0.4	0.3	0.2	0.2	0.2	0.0	
Without a needle ^k	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.6	0.4	0.6	0.4	0.4	0.7	+0.3	
Other Narcotics ^l	2.1	2.0	2.8	2.1	2.4	2.4	2.1	1.8	1.8	1.8	2.3	2.0	1.8	1.6	1.6	1.5	1.1	1.2	1.3	1.5	1.8	2.0	2.3	2.4	2.6	2.9	+0.3	
Amphetamines ^{b,l}	8.5	7.7	8.8	8.7	9.9	12.1	15.8	10.7	8.9	8.3	6.8	5.5	5.2	4.6	4.2	3.7	3.2	2.8	3.7	4.0	4.0	4.1	4.8	4.6	4.5	5.0	+0.5	
Methamphetamine ^m	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1.7	1.9	+0.2	
Crystal Meth. (Ice) ^m	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.6	0.6	0.5	0.6	0.7	1.1	1.1	0.8	1.2	0.8	1.0	+0.2	
Sedatives ^{l,n}	5.4	4.5	5.1	4.2	4.4	4.8	4.6	3.4	3.0	2.3	2.4	2.2	1.7	1.4	1.6	1.4	1.5	1.2	1.3	1.8	2.3	2.3	2.1	2.8	2.8	3.1	+0.3	
Barbiturates ^l	4.7	3.9	4.3	3.2	3.2	2.9	2.6	2.0	2.1	1.7	2.0	1.8	1.4	1.2	1.4	1.3	1.4	1.1	1.3	1.7	2.2	2.1	2.1	2.6	2.6	3.0	+0.4	
Methaqualone ^{l,n}	2.1	1.6	2.3	1.9	2.3	3.3	3.1	2.4	1.8	1.1	1.0	0.8	0.6	0.5	0.6	0.2	0.2	0.4	0.1	0.4	0.4	0.6	0.3	0.6	0.4	0.2	-0.3	
Tranquilizers ^l	4.1	4.0	4.6	3.4	3.7	3.1	2.7	2.4	2.5	2.1	2.1	2.1	2.0	1.5	1.3	1.2	1.4	1.0	1.2	1.4	1.8	2.0	1.8	2.4	2.5	2.6	+0.1	
Rohypnol ^e	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.5	0.3	0.3	0.3	0.4	+0.1	
Alcohol ^o	68.2	68.3	71.2	72.1	71.8	72.0	70.7	69.7	69.4	67.2	65.9	65.3	66.4	63.9	60.0	57.1	54.0	51.3	51.0	—	—	—	—	—	—	—	—	
Been Drunk ^m	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	31.6	29.9	28.9	30.8	33.2	31.3	34.2	32.9	32.9	32.3	-0.6	
Cigarettes	36.7	38.8	38.4	36.7	34.4	30.5	29.4	30.0	30.3	29.3	30.1	29.6	29.4	28.7	28.6	29.4	28.3	27.8	29.9	31.2	33.5	34.0	36.5	35.1	34.6	31.4	-3.2ss	
Smokeless Tobacco ^{e,p}	—	—	—	—	—	—	—	—	—	—	—	—	11.5	11.3	10.3	8.4	—	—	11.4	10.7	11.1	12.2	9.8	9.7	8.8	8.4	7.6	-0.7
Steroids ^m	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.8	1.0	0.8	0.6	0.7	0.9	0.7	0.7	1.0	1.1	0.9	0.8	-0.1	

NOTES: Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. See Table 4 for relevant footnotes. Any apparent inconsistency between the change estimate and the prevalence of use estimates for the two most recent classes is due to rounding error.

SOURCE: The Monitoring the Future Study, the University of Michigan.