

**TABLE 15**  
**Long-Term Trends in Annual Prevalence of Use of Various Drugs for Twelfth Graders**

	Percentage who used in last twelve months															
	Class of:															
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
Approx. N =	9,400	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Any Illicit Drug <sup>a,b</sup>	45.0	48.1	51.1	53.8	54.2	53.1	52.1	49.4	47.4	45.8	46.3	44.3	41.7	38.5	35.4	32.5
Any Illicit Drug Other Than Marijuana <sup>a,b,c</sup>	26.2	25.4	26.0	27.1	28.2	30.4	34.0	30.1	28.4	28.0	27.4	25.9	24.1	21.1	20.0	17.9
Marijuana/Hashish	40.0	44.5	47.6	50.2	50.8	48.8	46.1	44.3	42.3	40.0	40.6	38.8	36.3	33.1	29.6	27.0
Inhalants <sup>d</sup>	—	3.0	3.7	4.1	5.4	4.6	4.1	4.5	4.3	5.1	5.7	6.1	6.9	6.5	5.9	6.9
Inhalants, Adjusted <sup>d,e</sup>	—	—	—	—	8.9	7.9	6.1	6.6	6.2	7.2	7.5	8.9	8.1	7.1	6.9	7.5
Amyl/Butyl Nitrites <sup>f,g</sup>	—	—	—	—	6.5	5.7	3.7	3.6	3.6	4.0	4.0	4.7	2.6	1.7	1.7	1.4
Hallucinogens <sup>c</sup>	11.2	9.4	8.8	9.6	9.9	9.3	9.0	8.1	7.3	6.5	6.3	6.0	6.4	5.5	5.6	5.9
Hallucinogens, Adjusted <sup>c,h</sup>	—	—	—	—	11.8	10.4	10.1	9.0	8.3	7.3	7.6	7.6	6.7	5.8	6.2	6.0
LSD	7.2	6.4	5.5	6.3	6.6	6.5	6.5	6.1	5.4	4.7	4.4	4.5	5.2	4.8	4.9	5.4
Hallucinogens Other Than LSD <sup>c</sup>	9.4	7.0	6.9	7.3	6.8	6.2	5.6	4.7	4.1	3.8	3.6	3.0	3.2	2.1	2.2	2.1
PCP <sup>f,g</sup>	—	—	—	—	7.0	4.4	3.2	2.2	2.6	2.3	2.9	2.4	1.3	1.2	2.4	1.2
MDMA (Ecstasy) <sup>f</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Cocaine	5.6	6.0	7.2	9.0	12.0	12.3	12.4	11.5	11.4	11.6	13.1	12.7	10.3	7.9	6.5	5.3
Crack <sup>i</sup>	—	—	—	—	—	—	—	—	—	—	—	4.1	3.9	3.1	3.1	1.9
Other Cocaine <sup>j</sup>	—	—	—	—	—	—	—	—	—	—	—	—	9.8	7.4	5.2	4.6
Heroin <sup>k</sup>	1.0	0.8	0.8	0.8	0.5	0.5	0.5	0.6	0.6	0.5	0.6	0.5	0.5	0.5	0.6	0.5
With a needle <sup>l</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Without a needle <sup>l</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other Narcotics <sup>m,n</sup>	5.7	5.7	6.4	6.0	6.2	6.3	5.9	5.3	5.1	5.2	5.9	5.2	5.3	4.6	4.4	4.5
OxyContin <sup>o</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Vicodin <sup>o</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Amphetamines <sup>b,m</sup>	16.2	15.8	16.3	17.1	18.3	20.8	26.0†	20.3	17.9	17.7	15.8	13.4	12.2	10.9	10.8	9.1
Ritalin <sup>o</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Methamphetamine <sup>o</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Crystal Meth. (Ice) <sup>o</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1.3
Sedatives (Barbiturates) <sup>m</sup>	10.7	9.6	9.3	8.1	7.5	6.8	6.6	5.5	5.2	4.9	4.6	4.2	3.6	3.2	3.3	3.4
Sedatives, Adjusted <sup>m,p</sup>	11.7	10.7	10.8	9.9	9.9	10.3	10.5	9.1	7.9	6.6	5.8	5.2	4.1	3.7	3.7	3.6
Methaqualone <sup>m,q</sup>	5.1	4.7	5.2	4.9	5.9	7.2	7.6	6.8	5.4	3.8	2.8	2.1	1.5	1.3	1.3	0.7
Tranquilizers <sup>c,m</sup>	10.6	10.3	10.8	9.9	9.6	8.7	8.0	7.0	6.9	6.1	6.1	5.8	5.5	4.8	3.8	3.5
Rohypnol <sup>f</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
GHB <sup>t</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Ketamine <sup>t</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—

(Table continued on next page)

**TABLE 15 (cont'd)**  
**Long-Term Trends in Annual Prevalence of Use of Various Drugs for Twelfth Graders**

	Percentage who used in last twelve months															
	<u>Class of:</u>															
	<u>1975</u>	<u>1976</u>	<u>1977</u>	<u>1978</u>	<u>1979</u>	<u>1980</u>	<u>1981</u>	<u>1982</u>	<u>1983</u>	<u>1984</u>	<u>1985</u>	<u>1986</u>	<u>1987</u>	<u>1988</u>	<u>1989</u>	<u>1990</u>
<i>Approx. N =</i>	<i>9,400</i>	<i>15,400</i>	<i>17,100</i>	<i>17,800</i>	<i>15,500</i>	<i>15,900</i>	<i>17,500</i>	<i>17,700</i>	<i>16,300</i>	<i>15,900</i>	<i>16,000</i>	<i>15,200</i>	<i>16,300</i>	<i>16,300</i>	<i>16,700</i>	<i>15,200</i>
Alcohol	84.8	85.7	87.0	87.7	88.1	87.9	87.0	86.8	87.3	86.0	85.6	84.5	85.7	85.3	82.7	80.6
Been Drunk <sup>o</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Cigarettes	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Bidis <sup>o</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Kreteks <sup>o</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Smokeless Tobacco <sup>f,s</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Steroids <sup>o</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1.9	1.7

Cont'd 

NOTE: See Table 14 for relevant footnotes.

**TABLE 15 (cont'd)**  
**Long-Term Trends in Annual Prevalence of Use of Various Drugs for Twelfth Graders**

	Percentage who used in last twelve months															'04-'05 change
	Class of:															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	
<i>Approx. N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	
Any Illicit Drug <sup>a,b</sup>	29.4	27.1	31.0	35.8	39.0	40.2	42.4	41.4	42.1	40.9	41.4	41.0	39.3	38.8	38.4	-0.4
Any Illicit Drug Other Than Marijuana <sup>a,b,c</sup>	16.2	14.9	17.1	18.0	19.4	19.8	20.7	20.2	20.7	20.4†	21.6	20.9	19.8	20.5	19.7	-0.7
Marijuana/Hashish	23.9	21.9	26.0	30.7	34.7	35.8	38.5	37.5	37.8	36.5	37.0	36.2	34.9	34.3	33.6	-0.6
Inhalants <sup>d</sup>	6.6	6.2	7.0	7.7	8.0	7.6	6.7	6.2	5.6	5.9	4.5	4.5	3.9	4.2	5.0	+0.9
Inhalants, Adjusted <sup>d,e</sup>	6.9	6.4	7.4	8.2	8.4	8.5	7.3	7.1	6.0	6.2	4.9	4.9	4.5	4.6	5.4	+0.8
Amyl/Butyl Nitrites <sup>f,g</sup>	0.9	0.5	0.9	1.1	1.1	1.6	1.2	1.4	0.9	0.6	0.6	1.1	0.9	0.8	0.6	-0.2
Hallucinogens <sup>c</sup>	5.8	5.9	7.4	7.6	9.3	10.1	9.8	9.0	9.4	8.1‡	9.1	6.6	5.9	6.2	5.5	-0.7
Hallucinogens, Adjusted <sup>c,h</sup>	6.1	6.2	7.8	7.8	9.7	10.7	10.0	9.2	9.8	8.7‡	9.7	7.2	6.5	6.4	5.9	-0.4
LSD	5.2	5.6	6.8	6.9	8.4	8.8	8.4	7.6	8.1	6.6	6.6	3.5	1.9	2.2	1.8	-0.4
Hallucinogens Other Than LSD <sup>c</sup>	2.0	1.7	2.2	3.1	3.8	4.4	4.6	4.6	4.3	4.4‡	5.9	5.4	5.4	5.6	5.0	-0.6
PCP <sup>f,g</sup>	1.4	1.4	1.4	1.6	1.8	2.6	2.3	2.1	1.8	2.3	1.8	1.1	1.3	0.7	1.3	+0.6
MDMA (Ecstasy) <sup>f</sup>	—	—	—	—	—	4.6	4.0	3.6	5.6	8.2	9.2	7.4	4.5	4.0	3.0	-0.9
Cocaine	3.5	3.1	3.3	3.6	4.0	4.9	5.5	5.7	6.2	5.0	4.8	5.0	4.8	5.3	5.1	-0.2
Crack <sup>i</sup>	1.5	1.5	1.5	1.9	2.1	2.1	2.4	2.5	2.7	2.2	2.1	2.3	2.2	2.3	1.9	-0.3
Other Cocaine <sup>j</sup>	3.2	2.6	2.9	3.0	3.4	4.2	5.0	4.9	5.8	4.5	4.4	4.4	4.2	4.7	4.5	-0.2
Heroin <sup>k</sup>	0.4	0.6	0.5	0.6	1.1	1.0	1.2	1.0	1.1	1.5	0.9	1.0	0.8	0.9	0.8	0.0
With a needle <sup>l</sup>	—	—	—	—	0.5	0.5	0.5	0.4	0.4	0.4	0.3	0.4	0.4	0.4	0.5	+0.1
Without a needle <sup>l</sup>	—	—	—	—	1.0	1.0	1.2	0.8	1.0	1.6	0.8	0.8	0.8	0.7	0.8	+0.1
Other Narcotics <sup>m,n</sup>	3.5	3.3	3.6	3.8	4.7	5.4	6.2	6.3	6.7	7.0	6.7‡	9.4	9.3	9.5	9.0	-0.5
OxyContin <sup>o</sup>	—	—	—	—	—	—	—	—	—	—	—	4.0	4.5	5.0	5.5	+0.6
Vicodin <sup>o</sup>	—	—	—	—	—	—	—	—	—	—	—	9.6	10.5	9.3	9.5	+0.2
Amphetamines <sup>b,m</sup>	8.2	7.1	8.4	9.4	9.3	9.5	10.2	10.1	10.2	10.5	10.9	11.1	9.9	10.0	8.6	-1.4 s
Ritalin <sup>o</sup>	—	—	—	—	—	—	—	—	—	—	5.1	4.0	4.0	5.1	4.4	-0.7
Methamphetamine <sup>o</sup>	—	—	—	—	—	—	—	—	4.7	4.3	3.9	3.6	3.2	3.4	2.5	-0.9 s
Crystal Meth. (Ice) <sup>o</sup>	1.4	1.3	1.7	1.8	2.4	2.8	2.3	3.0	1.9	2.2	2.5	3.0	2.0	2.1	2.3	+0.3
Sedatives (Barbiturates) <sup>m</sup>	3.4	2.8	3.4	4.1	4.7	4.9	5.1	5.5	5.8	6.2	5.7	6.7	6.0	6.5	7.2	+0.7
Sedatives, Adjusted <sup>m,p</sup>	3.6	2.9	3.4	4.2	4.9	5.3	5.4	6.0	6.3	6.3	5.9	7.0	6.2	6.6	7.6	+0.9 s
Methaqualone <sup>m,q</sup>	0.5	0.6	0.2	0.8	0.7	1.1	1.0	1.1	1.1	0.3	0.8	0.9	0.6	0.8	0.9	+0.1
Tranquillizers <sup>c,m</sup>	3.6	2.8	3.5	3.7	4.4	4.6	4.7	5.5	5.8	5.7‡	6.9	7.7	6.7	7.3	6.8	-0.6
Rohypnol <sup>f</sup>	—	—	—	—	—	1.1	1.2	1.4	1.0	0.8	0.9‡	1.6	1.3	1.6	1.2	-0.4
GHB <sup>t</sup>	—	—	—	—	—	—	—	—	—	1.9	1.6	1.5	1.4	2.0	1.1	-0.9 s
Ketamine <sup>t</sup>	—	—	—	—	—	—	—	—	—	2.5	2.5	2.6	2.1	1.9	1.6	-0.3

(Table continued on next page)

**TABLE 15 (cont'd)**  
**Long-Term Trends in Annual Prevalence of Use of Various Drugs for Twelfth Graders**

	Percentage who used in last twelve months															'04-'05 change
	Class of:															
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	
<i>Approx. N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	
Alcohol <sup>f</sup>	77.7	76.8†	72.7	73.0	73.7	72.5	74.8	74.3	73.8	73.2	73.3	71.5	70.1	70.6	68.6	-2.1 s
Been Drunk <sup>o</sup>	52.7	50.3	49.6	51.7	52.5	51.9	53.2	52.0	53.2	51.8	53.2	50.4	48.0	51.8	47.7	-4.1 s
Cigarettes	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Bidis <sup>o</sup>	—	—	—	—	—	—	—	—	—	9.2	7.0	5.9	4.0	3.6	3.3	-0.3
Kreteks <sup>o</sup>	—	—	—	—	—	—	—	—	—	—	10.1	8.4	6.7	6.5	7.1	+0.6
Smokeless Tobacco <sup>f,s</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Steroids <sup>o</sup>	1.4	1.1	1.2	1.3	1.5	1.4	1.4	1.7	1.8	1.7	2.4	2.5	2.1	2.5	1.5	-1.1 sss

SOURCE: The Monitoring the Future Study, the University of Michigan.