TABLE 9
Trends in <u>Disapproval</u> of Drug Use in <u>Grade 8</u>

Do you disapprove of people who	Percentage who "disapprove" or "strongly disapprove" a 2010- 2011															2010-						
	<u>1991</u>	1992	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	2000	<u>2001</u>	2002	2003	2004	2005	2006	2007	2008	2009	<u>2010</u>	2011	change
Try marijuana once or twice	84.6	82.1	79.2	72.9	70.7	67.5	67.6	69.0	70.7	72.5	72.4	73.3	73.8	75.9	75.3	76.0	78.7	76.6	75.3	73.5	74.4	+0.9
Smoke marijuana occasionally	89.5	88.1	85.7	80.9	79.7	76.5	78.1	78.4	79.3	80.6	80.6	80.9	81.5	83.1	82.4	82.2	84.5	82.6	81.9	79.9	81.1	+1.2
Smoke marijuana regularly	92.1	90.8	88.9	85.3	85.1	82.8	84.6	84.5	84.5	85.3	84.5	85.3	85.7	86.8	86.3	86.1	87.7	86.8	85.9	84.3	85.7	+1.3
Try inhalants once or twice b	84.9	84.0	82.5	81.6	81.8	82.9	84.1	83.0	85.2	85.4	86.6	86.1	85.1	85.1	84.6	83.4	84.1	82.3	83.1	83.1	82.9	-0.2
Take inhalants regularly <sup>b</sup>	90.6	90.0	88.9	88.1	88.8	89.3	90.3	89.5	90.3	90.2	90.5	90.4	89.8	90.1	89.8	89.0	89.5	88.5	88.4	88.9	88.5	-0.3
Take LSD once or twice c	_	_	77.1	75.2	71.6	70.9	72.1	69.1	69.4	66.7	64.6	62.6	61.0	58.1	58.5	53.9	53.5	52.6	53.2	53.7	55.4	+1.7
Take LSD regularly <sup>c</sup>	_	_	79.8	78.4	75.8	75.3	76.3	72.5	72.5	69.3	67.0	65.5	63.5	60.5	60.7	55.8	55.6	54.7	55.7	55.8	57.6	+1.8
Try ecstasy (MDMA) once or twice <sup>d</sup>	_	_	_	_	_	_	_	_	_	_	69.0	74.3	77.7	76.3	75.0	66.7	65.7	63.5	62.3	62.4	64.2	+1.8
Take ecstasy (MDMA) occasionally <sup>d</sup>	_	_	_	_	_	_	_	_	_	_	73.6	78.6	81.3	79.4	77.9	69.8	68.3	66.5	65.7	65.9	67.5	+1.5
Try crack once or twice b	91.7	90.7	89.1	86.9	85.9	85.0	85.7	85.4	86.0	85.4	86.0	86.2	86.4	87.4	87.6	87.2	88.6	87.2	88.4	89.1	88.5	-0.7
Take crack occasionally <sup>b</sup>	93.3	92.5	91.7	89.9	89.8	89.3	90.3	89.5	89.9	88.8	89.8	89.6	89.8	90.3	90.5	90.0	91.2	90.3	91.0	91.5	91.0	-0.5
Try cocaine powder once or twice b	91.2	89.6	88.5	86.1	85.3	83.9	85.1	84.5	85.2	84.8	85.6	85.8	85.6	86.8	87.0	86.5	88.2	86.8	88.1	88.4	88.3	0.0
Take cocaine powder occasionally b	93.1	92.4	91.6	89.7	89.7	88.7	90.1	89.3	89.9	88.8	89.6	89.9	89.8	90.3	90.7	90.2	91.0	90.1	90.7	91.4	91.3	-0.2
Try heroin once or twice without using																						
a needle <sup>c</sup>	_	_	_	_	85.8	85.0	87.7	87.3	88.0	87.2	87.2	87.8	86.9	86.6	86.9	87.2	88.4	86.9	88.6	89.5	87.5	-2.0
Take heroin occasionally without using																						
a needle <sup>c</sup>	_	_	_	_	88.5	87.7	90.1	89.7	90.2	88.9	88.9	89.6	89.0	88.6	88.5	88.5	89.7	88.2	90.1	90.6	89.0	-1.6
Try one or two drinks of an alcoholic																						
beverage (beer, wine, liquor)	51.7	52.2	50.9	47.8	48.0	45.5	45.7	47.5	48.3	48.7	49.8	51.1	49.7	51.1	51.2	51.3	54.0	52.5	52.7	54.2	54.0	-0.2
Take one or two drinks nearly every day	82.2	81.0	79.6	76.7	75.9	74.1	76.6	76.9	77.0	77.8	77.4	78.3	77.1	78.6	78.7	78.7	80.4	79.2	78.5	79.5	80.7	+1.2
Have five or more drinks once or twice																						
each weekend	85.2	83.9	83.3	80.7	80.7	79.1	81.3	81.0	80.3	81.2	81.6	81.9	81.9	82.3	82.9	82.0	83.8	83.2	83.2	83.6	84.8	+1.2
Smoke one to five cigarettes per day <sup>d</sup>	_	_	_	_	_	_	_	_	75.1	79.1	80.4	81.1	81.4	83.1	82.9	83.5	85.3	85.0	83.6	84.7	86.8	+2.2
Smoke one or more packs of cigarettes																						
per day <sup>e</sup>	82.8	82.3	80.6	78.4	78.6	77.3	80.3	80.0	81.4	81.9	83.5	84.6	84.6	85.7	85.3	85.6	87.0	86.7	87.1	87.0	88.0	+1.0
Use smokeless tobacco regularly	79.1	77.2	77.1	75.1	74.0	74.1	76.5	76.3	78.0	79.2	79.4	80.6	80.7	81.0	82.0	81.0	82.3	82.1	81.5	81.2	82.6	+1.4
Take steroids <sup>f</sup>	89.8	90.3	89.9	87.9	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
Approximate weighted N =	: 17,400	18,500	18,400	17,400	17,600	18,000	18,800	18,100	16,700	16,700	16,200	15,100	16,500	17,000	16,800	16,500	16,100	15,700	15,000	15,300	16,000	)

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

<sup>&</sup>lt;sup>a</sup>Answer alternatives were: (1) Don't disapprove, (2) Disapprove, (3) Strongly disapprove, and (4) Can't say, drug unfamiliar. Percentages are shown for categories (2) and (3) combined.

 $<sup>^{\</sup>mathrm{b}}$ Beginning in 1997, data based on two thirds of N indicated due to changes in questionnaire forms.

<sup>&</sup>lt;sup>c</sup>Data based on one of two forms in 1993–1996; N is one half of N indicated. Beginning in 1997, data based on one third of N indicated due to changes in questionnaire forms.

<sup>&</sup>lt;sup>d</sup>Data based on one third of N indicated.

<sup>&</sup>lt;sup>e</sup>Beginning in 1999, data based on two thirds of *N* indicated due to changes in questionnaire forms.

<sup>&</sup>lt;sup>1</sup>Data based on two forms in 1991 and 1992. Data based on one of two forms in 1993 and 1994; N is one half of N indicated.