TABLE 6
Trends in <u>Harmfulness</u> of Drugs as Perceived by <u>8th Graders</u>

How much do you think people risk										Percer	ntage say	/ing grea	at risk <sup>a</sup>										2011-
harming themselves (physically or in other ways), if they	1991	<u>1992</u>	1993	1994	1995	<u>1996</u>	1997	1998	<u>1999</u>	2000	2001	<u>2002</u>	2003	2004	2005	2006	2007	2008	2009	<u>2010</u>	2011	<u>2012</u>	2012 change
Try marijuana once or twice <sup>b</sup>	40.4	39.1	36.2	31.6	28.9	27.9	25.3	28.1	28.0	29.0	27.7	28.2	30.2	31.9	31.4	32.2	32.8	31.1	29.5	29.5	28.2	26.0	-2.2 s
Smoke marijuana occasionally <sup>b</sup>	57.9	56.3	53.8	48.6	45.9	44.3	43.1	45.0	45.7	47.4	46.3	46.0	48.6	50.5	48.9	48.9	50.2	48.1	44.8	44.1	43.4	41.7	-1.7
Smoke marijuana regularly <sup>b</sup>	83.8	82.0	79.6	74.3	73.0	70.9	72.7	73.0	73.3	74.8	72.2	71.7	74.2	76.2	73.9	73.2	74.3	72.0	69.8	68.0	68.3	66.9	-1.5
	03.0	02.0	79.0	74.3	73.0	70.9	12.1	73.0	13.3	74.0	12.2	11.1	74.2	70.2	13.9	13.2	74.5	72.0	09.0	00.0	00.3	24.4	-1.5
Try synthetic marijuana once or twice <sup>c</sup> Take synthetic marijuana occasionally <sup>c</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	24.4 36.8	_
Try inhalants once or twice <sup>d</sup>					-	-	-		-		-	-	-										_
	35.9	37.0	36.5	37.9	36.4	40.8	40.1	38.9	40.8	41.2	45.6	42.8	40.3	38.7	37.5	35.8	35.9	33.9	34.1	35.5	34.7	34.2	-0.6
Take inhalants regularly d	65.6	64.4	64.6	65.5	64.8	68.2	68.7	67.2	68.8	69.9	71.6	69.9	67.4	66.4	64.1	62.1	61.9	59.2	58.1	60.6	59.0	59.0	0.0
Take LSD once or twice <sup>e</sup>	_	_	42.1	38.3	36.7	36.5	37.0	34.9	34.1	34.0	31.6	29.6	27.9	26.8	25.8	23.8	22.8	21.9	21.4	23.6	21.7	19.9	-1.9
Take LSD regularly <sup>e</sup>	-	-	68.3	65.8	64.4	63.6	64.1	59.6	58.8	57.5	52.9	49.3	48.2	45.2	44.0	40.0	38.5	36.9	37.0	38.6	37.8	35.0	-2.8
Try ecstasy (MDMA) once or twice c	_	_	_	_	_	_	_	_	_	_	35.8	38.9	41.9	42.5	40.0	32.8	30.4	28.6	26.0	27.0	25.4	23.6	-1.8
Take ecstasy (MDMA) occasionally <sup>c</sup>	—	—	—	—	—	—	—	-	—	-	55.5	61.8	65.8	65.1	60.8	52.0	48.6	46.8	43.9	45.0	43.7	41.0	-2.7
Try salvia once or twice c	-	-	-	-	-	-	-	-	-	-	-	-	-	—	-	-	-	-	-	-	-	9.5	-
Take salvia occasionally °	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	16.1	-
Try crack once or twice d	62.8	61.2	57.2	54.4	50.8	51.0	49.9	49.3	48.7	48.5	48.6	47.4	48.7	49.0	49.6	47.6	47.3	47.1	46.6	49.6	48.1	47.0	-1.2
Take crack occasionally <sup>d</sup>	82.2	79.6	76.8	74.4	72.1	71.6	71.2	70.6	70.6	70.1	70.0	69.7	70.3	70.4	69.4	68.7	68.3	67.9	66.6	68.4	67.7	67.8	+0.1
Try cocaine powder once or twice <sup>d</sup>	55.5	54.1	50.7	48.4	44.9	45.2	45.0	44.0	43.3	43.3	43.9	43.2	43.7	44.4	44.2	43.5	43.5	42.7	42.3	45.7	43.3	42.8	-0.5
Take cocaine powder occasionally <sup>d</sup>	77.0	74.3	71.8	69.1	66.4	65.7	65.8	65.2	65.4	65.5	65.8	64.9	65.8	66.0	65.3	64.0	64.2	62.7	62.3	64.2	63.5	63.3	-0.2
Try heroin once or twice without using a needle <sup>e</sup>	_	_	_	_	60.1	61.3	63.0	62.8	63.0	62.0	61.1	62.6	62.7	61.6	61.4	60.4	60.3	60.8	60.0	62.3	61.7	59.1	-2.6
Take heroin occasionally without using																							
a needle <sup>e</sup>	_	_	_	_	76.8	76.6	79.2	79.0	78.9	78.6	78.5	78.5	77.8	77.5	76.8	75.3	76.4	75.5	74.0	76.7	75.9	75.1	-0.8
Try OxyContin once or twice <sup>c</sup>	-	-	_	-	-	-	-	-	-	-	-	-	-	-	-	-	-	_	_	-	_	21.9	_
Take OxyContin occasionally <sup>c</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	35.3	_
Try Vicodin once or twice <sup>c</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	17.5	_
Take Vicodin occasionally <sup>c</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	29.4	_
Try Adderall once or twice <sup>c</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	17.6	_
Take Adderall occasionally <sup>c</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	29.9	_
Try bath salts (synthetic stimulants)																							
once or twice <sup>c</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	24.9	_
Take bath salts (synthetic stimulants)																							
occasionally <sup>c</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	38.8	_
Try cough/cold medicine once or twice <sup>c</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	21.2	_
Take cough/cold medicine occasionally <sup>c</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	38.8	_
Try one or two drinks of an alcoholic	-	_	_	-	-	-	-	-	-	-	_	_	_	_	-	_	-	_	_	_	-	00.0	
beverage (beer, wine, liquor) <sup>b</sup>	11.0	12.1	12.4	11.6	11.6	11.8	10.4	12.1	11.6	11.9	12.2	12.5	12.6	13.7	13.9	14.2	14.9	13.5	14.4	14.9	14.5	13.9	-0.7
Take one or two drinks nearly every day <sup>b</sup>	31.8	32.4	32.6	29.9	30.5	28.6	29.1	30.3	29.7	30.4	30.0	29.6	29.9	31.0	31.4	31.3	32.6	31.5	31.5	32.3	31.8	31.4	-0.7
	31.0	32.4	32.0	29.9	30.5	20.0	29.1	30.3	29.7	30.4	30.0	29.0	29.9	31.0	31.4	31.3	32.0	31.5	31.5	32.3	31.0	31.4	-0.4
Have five or more drinks once or twice	50.4					54.0							50.5	50.0									0.0
each weekend <sup>b</sup>	59.1	58.0	57.7	54.7	54.1	51.8	55.6	56.0	55.3	55.9	56.1	56.4	56.5	56.9	57.2	56.4	57.9	57.0	55.8	57.2	58.4	58.2	-0.3
Smoke one to five cigarettes per day <sup>c</sup>	-	-	-	-	-	-	-	-	26.9	28.9	30.5	32.8	33.4	37.0	37.5	37.0	38.6	38.6	38.6	38.2	37.4	40.4	+3.0
Smoke one or more packs of cigarettes																							
per day <sup>†</sup>	51.6	50.8	52.7	50.8	49.8	50.4	52.6	54.3	54.8	58.8	57.1	57.5	57.7	62.4	61.5	59.4	61.1	59.8	59.1	60.9	62.5	62.6	+0.1
Use smokeless tobacco regularly	35.1	35.1	36.9	35.5	33.5	34.0	35.2	36.5	37.1	39.0	38.2	39.4	39.7	41.3	40.8	39.5	41.8	41.0	40.8	41.8	40.8	37.8	-2.9 s
Take dissolvable tobacco regularly °	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	34.8	-
Take snus regularly <sup>c</sup>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	42.2	-
Take steroids <sup>g</sup>	64.2	69.5	70.2	67.6	-	-	-	-	-	—	-	-	-	-	-	-	-	-	-	-	-	-	_
Approximate weighted N =	17,400	18,700	18,400	17,400	17,500	17,900	18,800	18,100	16,700	16,700	16,200	15,100	16,500	17,000	16,800	16,500	16,100	15,700	15,000	15,300	16,000	15,100	

Approximate weighted N = 17,400 18,700 18,400 17,400 17,500 17,900 18,800 18,100 16,700 16,700 16,200 15,100 16,500 17,000 16,800 16,500 15,100 15,700 15,000 15,300 16,000 15,100

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' - ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates

for the two most recent years is due to rounding.

<sup>a</sup>Answer alternatives were: (1) No risk, (2) Slight risk, (3) Moderate risk, (4) Great risk, and (5) Can't say, drug unfamiliar.

<sup>b</sup>Beginning in 2012 data based on two thirds of N indicated.

<sup>c</sup>Data based on one third of *N* indicated.

<sup>d</sup>Beginning in 1997, data based on two thirds of *N* indicated due to changes in questionnaire forms.

<sup>e</sup>Data based on one of two forms in 1993–1996; N is one half of N indicated. Beginning in 1997, data based on one third of N indicated due to changes in questionnaire forms.

<sup>1</sup>Beginning in 1999, data based on two thirds of *N* indicated due to changes in questionnaire forms.

<sup>9</sup>Data based on two forms in 1991 and 1992. Data based on one of two forms in 1993 and 1994; N is one half of N indicated.