

TABLE 9
Trends in Disapproval of Drug Use in Grade 10

Do you disapprove of people who . . .	Percentage who disapprove or strongly disapprove ^a														
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Try marijuana once or twice ^b	74.6	74.8	70.3	62.4	59.8	55.5	54.1	56.0	56.2	54.9	54.8	57.8	58.1	60.4	61.3
Smoke marijuana occasionally ^b	83.7	83.6	79.4	72.3	70.0	66.9	66.2	67.3	68.2	67.2	66.2	68.3	68.4	70.8	71.9
Smoke marijuana regularly ^b	90.4	90.0	87.4	82.2	81.1	79.7	79.7	80.1	79.8	79.1	78.0	78.6	78.8	81.3	82.0
Try inhalants once or twice ^c	85.2	85.6	84.8	84.9	84.5	86.0	86.9	85.6	88.4	87.5	87.8	88.6	87.7	88.5	88.1
Take inhalants regularly ^c	91.0	91.5	90.9	91.0	90.9	91.7	91.7	91.1	92.4	91.8	91.3	91.8	91.0	92.3	91.9
Take LSD once or twice ^d	—	—	82.1	79.3	77.9	76.8	76.6	76.7	77.8	77.0	75.4	74.6	74.4	72.4	71.8
Take LSD regularly ^d	—	—	86.8	85.6	84.8	84.5	83.4	82.9	84.3	82.1	80.8	79.4	77.6	75.9	75.0
Try ecstasy (MDMA) once or twice ^e	—	—	—	—	—	—	—	—	—	—	72.6	77.4	81.0	83.7	83.1
Take ecstasy (MDMA) occasionally ^e	—	—	—	—	—	—	—	—	—	—	81.0	84.6	86.3	88.0	87.4
Try crack once or twice ^c	92.5	92.5	91.4	89.9	88.7	88.2	87.4	87.1	87.8	87.1	86.9	88.0	87.6	88.6	88.8
Take crack occasionally ^c	94.3	94.4	93.6	92.5	91.7	91.9	91.0	90.6	91.5	90.9	90.6	91.0	91.0	91.8	91.8
Try cocaine powder once or twice ^c	90.8	91.1	90.0	88.1	86.8	86.1	85.1	84.9	86.0	84.8	85.3	86.4	85.9	86.8	86.9
Take cocaine powder occasionally ^c	94.0	94.0	93.2	92.1	91.4	91.1	90.4	89.7	90.7	89.9	90.2	89.9	90.4	91.2	91.2
Try heroin once or twice without using a needle ^d	—	—	—	—	89.7	89.5	89.1	88.6	90.1	90.1	89.1	89.2	89.3	90.1	90.3
Take heroin occasionally without using a needle ^d	—	—	—	—	91.6	91.7	91.4	90.5	91.8	92.3	90.8	90.7	90.6	91.8	92.0
Try one or two drinks of an alcoholic beverage (beer, wine, liquor) ^b	37.6	39.9	38.5	36.5	36.1	34.2	33.7	34.7	35.1	33.4	34.7	37.7	36.8	37.6	38.5
Take one or two drinks nearly every day ^b	81.7	81.7	78.6	75.2	75.4	73.8	75.4	74.6	75.4	73.8	73.8	74.9	74.2	75.1	76.9
Have five or more drinks once or twice each weekend ^b	76.7	77.6	74.7	72.3	72.2	70.7	70.2	70.5	69.9	68.2	69.2	71.5	71.6	71.8	73.7
Smoke one to five cigarettes per day ^e	—	—	—	—	—	—	—	—	67.8	69.1	71.2	74.3	76.2	77.5	79.3
Smoke one or more packs of cigarettes per day ^f	79.4	77.8	76.5	73.9	73.2	71.6	73.8	75.3	76.1	76.7	78.2	80.6	81.4	82.7	84.3
Use electronic cigarettes (e-cigarettes) regularly ^e	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Vape an e-liquid with nicotine occasionally ^e	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Vape an e-liquid with nicotine regularly ^e	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Use smokeless tobacco regularly ^b	75.4	74.6	73.8	71.2	71.0	71.0	72.3	73.2	75.1	75.8	76.1	78.7	79.4	80.2	80.5
Take steroids ^g	90.0	91.0	91.2	90.8	—	—	—	—	—	—	—	—	—	—	—

Table continued on next page.

Approximate weighted N = 14,800 14,800 15,300 15,900 17,000 15,700 15,600 15,000 13,600 14,300 14,000 14,300 15,800 16,400 16,200

TABLE 9 (cont.)
Trends in Disapproval of Drug Use in Grade 10

Do you disapprove of people who . . .	Percentage who disapprove or strongly disapprove ^a												2016–2017 change
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	
Try marijuana once or twice ^b	62.5	63.9	64.5	60.1	59.2	58.5	56.2	53.2	53.8	52.7	52.6	48.1	-4.6 sss
Smoke marijuana occasionally ^b	72.6	73.3	73.6	69.2	68.0	67.9	65.7	62.1	62.9	62.6	61.9	58.1	-3.8 ss
Smoke marijuana regularly ^b	82.5	82.4	83.0	79.9	78.7	78.8	77.3	73.8	74.6	74.3	73.5	70.2	-3.3 ss
Try inhalants once or twice ^c	88.1	87.6	87.1	87.0	86.5	86.9	85.7	86.1	85.9	84.1	83.3	80.7	-2.6 s
Take inhalants regularly ^c	92.2	91.8	91.6	91.1	90.8	90.9	90.0	89.7	89.7	88.3	87.1	85.4	-1.8
Take LSD once or twice ^d	71.2	67.7	66.3	67.8	68.2	68.5	68.3	69.1	67.8	70.3	69.5	66.9	-2.7
Take LSD regularly ^d	74.9	71.5	69.8	72.2	72.9	72.5	73.0	74.2	73.3	76.5	74.9	74.5	-0.4
Try ecstasy (MDMA) once or twice ^e	81.6	80.0	78.1	76.5	75.5	76.1	75.3	75.4	74.4‡	78.0	76.8	74.7	-2.1
Take ecstasy (MDMA) occasionally ^e	86.0	84.3	83.0	81.3	81.3	82.2	81.2	81.3	80.4‡	84.0	81.7	80.0	-1.7
Try crack once or twice ^c	89.5	89.5	90.8	90.4	90.3	90.9	91.0	90.6	90.6	90.1	89.7	88.4	-1.3
Take crack occasionally ^c	92.0	92.7	92.9	92.8	92.4	93.0	93.0	92.4	92.4	92.1	91.1	90.0	-1.0
Try cocaine powder once or twice ^c	87.3	87.7	88.6	88.4	89.0	89.4	89.3	88.7	88.9	87.9	87.9	86.1	-1.8 s
Take cocaine powder occasionally ^c	91.4	92.0	92.1	92.1	92.2	92.5	92.4	91.8	91.9	91.8	90.8	89.9	-0.9
Try heroin once or twice without using a needle ^d	91.1	90.7	91.4	91.6	91.4	91.6	91.9	91.3	91.9	91.7	90.2	89.7	-0.5
Take heroin occasionally without using a needle ^d	92.5	92.5	92.5	93.0	92.4	92.4	92.9	92.3	92.7	92.7	90.9	90.5	-0.5
Try one or two drinks of an alcoholic beverage (beer, wine, liquor) ^b	37.8	39.5	41.8	39.7	40.3	41.5	39.6	38.5	40.7	40.0	41.8	39.3	-2.5 s
Take one or two drinks nearly every day ^b	76.4	77.1	79.1	77.6	77.6	80.0	78.0	77.1	77.9	78.2	78.6	77.7	-0.9
Have five or more drinks once or twice each weekend ^b	72.9	74.1	77.2	75.1	75.9	77.3	77.5	77.8	79.5	79.6	80.8	80.1	-0.7
Smoke one to five cigarettes per day ^e	80.2	79.7	82.5	80.0	80.6	82.1	—	—	—	—	—	—	—
Smoke one or more packs of cigarettes per day ^f	83.2	84.7	85.2	84.5	83.9	85.8	86.0	86.1	88.0	88.3	88.5	87.8	-0.8
Use electronic cigarettes (e-cigarettes) regularly ^e	—	—	—	—	—	—	—	—	54.6	59.9	65.0	—	—
Vape an e-liquid with nicotine occasionally ^e	—	—	—	—	—	—	—	—	—	—	—	64.2	—
Vape an e-liquid with nicotine regularly ^e	—	—	—	—	—	—	—	—	—	—	—	73.9	—
Use smokeless tobacco regularly ^b	80.5	80.9	81.8	79.5	78.5	79.5	79.5	77.7	78.7	80.1	81.2	80.7	-0.5
Take steroids ^g	—	—	—	—	—	—	—	—	—	—	—	—	—

Table continued on next page.

Approximate weighted N = 16,200 16,100 15,100 15,900 15,200 14,900 15,000 12,900 13,000 15,600 14,700 13,500

TABLE 9 (cont.)
Trends in Disapproval of Drug Use in Grade 10

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: $s = .05$, $ss = .01$, $sss = .001$. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. '‡' indicates that the question changed the following year.

^aAnswer alternatives were: (1) Don't disapprove, (2) Disapprove, (3) Strongly disapprove, and (4) Can't say, drug unfamiliar. Percentages are shown for categories (2) and (3) combined.

^bBeginning in 2012, data based on two thirds of N indicated.

^cBeginning in 1997, data based on two thirds of N indicated due to changes in questionnaire forms.

^dData based on one of two forms in 1993–1996; N is one half of N indicated. Beginning in 1997, data based on one third of N indicated due to changes in questionnaire forms.

^eData based on one third of N indicated. For MDMA "Molly" was added to the question text in 2015; 2014 and 2015 data are not comparable due to this change.

^fBeginning in 1999, data based on two thirds of N indicated due to changes in questionnaire forms.

^gData based on two forms in 1991 and 1992. Data based on one of two forms in 1993 and 1994; N is one half of N indicated.