## TABLE 14Trends in <a href="mailto:Disapproval">Disapproval</a> of Drug Use in <a href="mailto:Grade 12">Grade 12</a>

Percentage who disapprove or strongly disapprove<sup>b</sup>

Do you disapprove of people (who are 18 or older) doing each of the following? <sup>a</sup>	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	<u>1990</u>	
Trying marijuana once or twice	47.0	38.4	33.4	33.4	34.2	39.0	40.0	45.5	46.3	49.3	51.4	54.6	56.6	60.8	64.6	67.8	
Smoking marijuana occasionally	54.8	47.8	44.3	43.5	45.3	49.7	52.6	59.1	60.7	63.5	65.8	69.0	71.6	74.0	77.2	80.5	
Smoking marijuana regularly	71.9	69.5	65.5	67.5	69.2	74.6	77.4	80.6	82.5	84.7	85.5	86.6	89.2	89.3	89.8	91.0	
Trying LSD once or twice	82.8	84.6	83.9	85.4	86.6	87.3	86.4	88.8	89.1	88.9	89.5	89.2	91.6	89.8	89.7	89.8	
Taking LSD regularly	94.1	95.3	95.8	96.4	96.9	96.7	96.8	96.7	97.0	96.8	97.0	96.6	97.8	96.4	96.4	96.3	
Trying ecstasy (MDMA, Molly) once or twice <sup>c</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
Trying cocaine once or twice	81.3	82.4	79.1	77.0	74.7	76.3	74.6	76.6	77.0	79.7	79.3	80.2	87.3	89.1	90.5	91.5	
Taking cocaine regularly	93.3	93.9	92.1	91.9	90.8	91.1	90.7	91.5	93.2	94.5	93.8	94.3	96.7	96.2	96.4	96.7	
Trying crack once or twice	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	92.3	
Taking crack occasionally	_	_	_	_	_	_	_	_	_	_	_		_	_	_	94.3	
Taking crack regularly	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	94.9	
Trying cocaine powder once or twice	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	87.9	
Taking cocaine powder occasionally	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	92.1	
Taking cocaine powder regularly	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	93.7	Table continued on next page
Trying heroin once or twice	91.5	92.6	92.5	92.0	93.4	93.5	93.5	94.6	94.3	94.0	94.0	93.3	96.2	95.0	95.4	95.1	
Taking heroin occasionally	94.8	96.0	96.0	96.4	96.8	96.7	97.2	96.9	96.9	97.1	96.8	96.6	97.9	96.9	97.2	96.7	
Taking heroin regularly	96.7	97.5	97.2	97.8	97.9	97.6	97.8	97.5	97.7	98.0	97.6	97.6	98.1	97.2	97.4	97.5	
Trying heroin once or twice without using a needle	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
Taking heroin occasionally without using a needle	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
Trying amphetamines once or twice <sup>d</sup>	74.8	75.1	74.2	74.8	75.1	75.4	71.1	72.6	72.3	72.8	74.9	76.5	80.7	82.5	83.3	85.3	
Taking amphetamines regularly <sup>d</sup>	92.1	92.8	92.5	93.5	94.4	93.0	91.7	92.0	92.6	93.6	93.3	93.5	95.4	94.2	94.2	95.5	
Trying sedatives (barbiturates) once or twice <sup>e</sup>	77.7	81.3	81.1	82.4	84.0	83.9	82.4	84.4	83.1	84.1	84.9	86.8	89.6	89.4	89.3	90.5	
Taking sedatives (barbiturates) regularly <sup>e</sup>	93.3	93.6	93.0	94.3	95.2	95.4	94.2	94.4	95.1	95.1	95.5	94.9	96.4	95.3	95.3	96.4	
Trying one or two drinks of an alcoholic beverage																	
(beer, wine, liquor)	21.6	18.2	15.6	15.6	15.8	16.0	17.2	18.2	18.4	17.4	20.3	20.9	21.4	22.6	27.3	29.4	
Taking one or two drinks nearly every day	67.6	68.9	66.8	67.7	68.3	69.0	69.1	69.9	68.9	72.9	70.9	72.8	74.2	75.0	76.5	77.9	
Taking four or five drinks nearly every day	88.7	90.7	88.4	90.2	91.7	90.8	91.8	90.9	90.0	91.0	92.0	91.4	92.2	92.8	91.6	91.9	
Having five or more drinks once or twice																	
each weekend	60.3	58.6	57.4	56.2	56.7	55.6	55.5	58.8	56.6	59.6	60.4	62.4	62.0	65.3	66.5	68.9	
Smoking one or more packs of cigarettes per day	67.5	65.9	66.4	67.0	70.3	70.8	69.9	69.4	70.8	73.0	72.3	75.4	74.3	73.1	72.4	72.8	
Vape an e-liquid with nicotine ocasionally $\int$	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
Vape an e-liquid with nicotine regularly <sup>f</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
Use JUUL occasionally <sup>f</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
Use JUUL regularly <sup>f</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
Taking steroids	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	90.8	
Approximate weighted N =	2.677	2,957	3,085	3,686	3,221	3,261	3.610	3,651	3,341	3,254	3,265	3,113	3,302	3,311	2,799	2,566	

## TABLE 14 (cont.)Trends in <u>Disapproval</u> of Drug Use in <u>Grade 12</u>

Percentage who disapprove or strongly disapprove<sup>b</sup>

Do you disapprove of people (who are 18 or older)																	
doing each of the following? <sup>a</sup>	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>	<u>2006</u>	
Trying marijuana once or twice	68.7	69.9	63.3	57.6	56.7	52.5	51.0	51.6	48.8	52.5	49.1	51.6	53.4	52.7	55.0	55.6	
Smoking marijuana occasionally	79.4	79.7	75.5	68.9	66.7	62.9	63.2	64.4	62.5	65.8	63.2	63.4	64.2	65.4	67.8	69.3	
Smoking marijuana regularly	89.3	90.1	87.6	82.3	81.9	80.0	78.8	81.2	78.6	79.7	79.3	78.3	78.7	80.7	82.0	82.2	
Trying LSD once or twice	90.1	88.1	85.9	82.5	81.1	79.6	80.5	82.1	83.0	82.4	81.8	84.6	85.5	87.9	87.9	88.0	
Taking LSD regularly	96.4	95.5	95.8	94.3	92.5	93.2	92.9	93.5	94.3	94.2	94.0	94.0	94.4	94.6	95.6	95.9	
Trying ecstasy (MDMA, Molly) once or twice <sup>c</sup>	_	_	—	_	_	_	82.2	82.5	82.1	81.0	79.5	83.6	84.7	87.7	88.4	89.0	
Trying cocaine once or twice	93.6	93.0	92.7	91.6	90.3	90.0	88.0	89.5	89.1	88.2	88.1	89.0	89.3	88.6	88.9	89.1	
Taking cocaine regularly	97.3	96.9	97.5	96.6	96.1	95.6	96.0	95.6	94.9	95.5	94.9	95.0	95.8	95.4	96.0	96.1	
Trying crack once or twice	92.1	93.1	89.9	89.5	91.4	87.4	87.0	86.7	87.6	87.5	87.0	87.8	86.6	86.9	86.7	88.8	
Taking crack occasionally	94.2	95.0	92.8	92.8	94.0	91.2	91.3	90.9	92.3	91.9	91.6	91.5	90.8	92.1	91.9	92.9	
Taking crack regularly	95.0	95.5	93.4	93.1	94.1	93.0	92.3	91.9	93.2	92.8	92.2	92.4	91.2	93.1	92.1	93.8	
Trying cocaine powder once or twice	88.0	89.4	86.6	87.1	88.3	83.1	83.0	83.1	84.3	84.1	83.3	83.8	83.6	82.2	83.2	84.1	
Taking cocaine powder occasionally	93.0	93.4	91.2	91.0	92.7	89.7	89.3	88.7	90.0	90.3	89.8	90.2	88.9	90.0	89.4	90.4	
Taking cocaine powder regularly	94.4	94.3	93.0	92.5	93.8	92.9	91.5	91.1	92.3	92.6	92.5	92.2	90.7	92.6	92.0	93.2	
Trying heroin once or twice	96.0	94.9	94.4	93.2	92.8	92.1	92.3	93.7	93.5	93.0	93.1	94.1	94.1	94.2	94.3	93.8	
Taking heroin occasionally	97.3	96.8	97.0	96.2	95.7	95.0	95.4	96.1	95.7	96.0	95.4	95.6	95.9	96.4	96.3	96.2	
Taking heroin regularly	97.8	97.2	97.5	97.1	96.4	96.3	96.4	96.6	96.4	96.6	96.2	96.2	97.1	97.1	96.7	96.9	
Trying heroin once or twice without using a needle	_	_	_	_	92.9	90.8	92.3	93.0	92.6	94.0	91.7	93.1	92.2	93.1	93.2	93.7	
Taking heroin occasionally without using a needle	_	_	_	_	94.7	93.2	94.4	94.3	93.8	95.2	93.5	94.4	93.5	94.4	95.0	94.5	
Trying amphetamines once or twice <sup>d</sup>	86.5	86.9	84.2	81.3	82.2	79.9	81.3	82.5	81.9	82.1	82.3	83.8	85.8	84.1	86.1	86.3	
Taking amphetamines regularly <sup>d</sup>	96.0	95.6	96.0	94.1	94.3	93.5	94.3	94.0	93.7	94.1	93.4	93.5	94.0	93.9	94.8	95.3	
Trying sedatives (barbiturates) once or twice <sup>e</sup>	90.6	90.3	89.7	87.5	87.3	84.9	86.4	86.0	86.6	85.9	85.9	86.6	87.8‡	83.7	85.4	85.3	
Taking sedatives (barbiturates) regularly <sup>e</sup>	97.1	96.5	97.0	96.1	95.2	94.8	95.3	94.6	94.7	95.2	94.5	94.7	94.4‡	94.2	95.2	95.1	
Trying one or two drinks of an alcoholic beverage																	
(beer, wine, liquor)	29.8	33.0	30.1	28.4	27.3	26.5	26.1	24.5	24.6	25.2	26.6	26.3	27.2	26.0	26.4	29.0	
Taking one or two drinks nearly every day	76.5	75.9	77.8	73.1	73.3	70.8	70.0	69.4	67.2	70.0	69.2	69.1	68.9	69.5	70.8	72.8	
Taking four or five drinks nearly every day	90.6	90.8	90.6	89.8	88.8	89.4	88.6	86.7	86.9	88.4	86.4	87.5	86.3	87.8	89.4	90.6	
Having five or more drinks once or twice																	
each weekend	67.4	70.7	70.1	65.1	66.7	64.7	65.0	63.8	62.7	65.2	62.9	64.7	64.2	65.7	66.5	68.5	
Smoking one or more packs of cigarettes per day	71.4	73.5	70.6	69.8	68.2	67.2	67.1	68.8	69.5	70.1	71.6	73.6	74.8	76.2	79.8	81.5	
Vape an e-liquid with nicotine ocasionally	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
Vape an e-liquid with nicotine regularly <sup>f</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
Use JUUL occasionally <sup>f</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
Use JUUL regularly <sup>f</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
Taking steroids	90.5	92.1	92.1	91.9	91.0	91.7	91.4	90.8	88.9	88.8	86.4	86.8	86.0	87.9	88.8	89.4	
Approximate weighted N =	2,547	2,645	2,723	2,588	2,603	2,399	2,601	2,545	2,310	2,150	2,144	2,160	2,442	2,455	2,460	2,377	

able continued on next page.

TABLE 14 (cont.)	
Trends in <b>Disapproval</b> of Drug Use in G	Frade 12

Percentage who disapprove or strongly disapprove<sup>b</sup>

Do you disapprove of people (who are 18 or older)														2018–2019
doing each of the following? <sup>a</sup>	<u>2007</u>	<u>2008</u>	<u>2009</u>	<u>2010</u>	<u>2011</u>	<u>2012</u>	<u>2013</u>	<u>2014</u>	<u>2015</u>	<u>2016</u>	<u>2017</u>	<u>2018</u>	<u>2019</u>	<u>change</u>
Trying marijuana once or twice	58.6	55.5	54.8	51.6	51.3	48.8	49.1	48.0	45.5	43.1	39.0	41.1	37.0	-4.2
Smoking marijuana occasionally	70.2	67.3	65.6	62.0	60.9	59.1	58.9	56.7	52.9	50.5	46.7	49.2	44.1	-5.1 s
Smoking marijuana regularly	83.3	79.6	80.3	77.7	77.5	77.8	74.5	73.4	70.7	68.5	64.7	66.7	65.1	-1.6
Trying LSD once or twice	87.8	85.5	88.2	86.5	86.3	87.2	86.6	85.0	81.7	82.4	78.0	80.5	76.9	-3.6 ss
Taking LSD regularly	94.9	93.5	95.3	94.3	94.9	95.2	95.3	94.7	92.5	92.4	92.7	93.4	93.3	-0.1
Trying ecstasy (MDMA, Molly) once or twice <sup>c</sup>	87.8	88.2	88.2	86.3	83.9	87.1	84.9‡	83.1	84.5	84.0	85.1	85.6	88.7	+3.1 s
Trying cocaine once or twice	89.6	89.2	90.8	90.5	91.1	91.0	92.3	90.0	89.0	88.4	88.0	88.9	88.7	-0.2
Taking cocaine regularly	96.2	94.8	96.5	96.0	96.0	96.8	96.7	96.3	95.2	94.8	94.8	95.8	96.1	+0.3
Trying crack once or twice	88.8	89.6	90.9	89.8	91.4	92.8	91.4	89.3	90.2	90.1	89.7	90.4	87.1	-3.4 s
Taking crack occasionally	92.4	93.3	94.0	92.6	93.9	95.0	93.6	91.9	92.5	92.0	91.8	92.2	89.0	-3.1 s
Taking crack regularly	93.6	93.5	94.3	93.1	94.4	95.4	94.1	92.4	92.8	92.6	92.5	92.5	89.4	-3.1 s
Trying cocaine powder once or twice	83.5	85.7	87.3	87.0	88.1	88.7	88.2	85.5	86.4	86.6	85.5	86.5	83.8	-2.7
Taking cocaine powder occasionally	90.6	91.7	92.3	91.0	92.2	93.0	91.7	90.4	91.3	90.6	90.3	91.3	88.1	-3.1 s
Taking cocaine powder regularly	92.6	92.8	93.9	92.6	93.8	95.0	94.1	91.7	92.4	92.0	92.2	92.0	89.4	-2.6
Trying heroin once or twice	94.8	93.3	94.7	93.9	94.3	95.8	95.6	94.7	94.2	94.1	93.7	95.0	94.7	-0.2
Taking heroin occasionally	96.8	95.3	96.9	96.2	96.3	97.0	96.9	96.6	95.3	95.5	95.5	96.4	96.3	0.0
Taking heroin regularly	97.1	95.9	97.4	96.4	96.7	97.4	97.4	97.1	96.4	95.7	95.9	96.8	96.8	0.0
Trying heroin once or twice without using a needle	93.6	94.2	94.7	93.2	92.6	95.2	93.7	92.5	92.6	93.8	93.3	93.0	95.1	+2.1 s
Taking heroin occasionally without using a needle	94.9	95.3	95.5	94.5	94.1	95.9	94.6	93.5	92.8	94.0	93.8	93.4	95.2	+1.8 s
Trying amphetamines once or twice <sup>d</sup>	87.3	87.2	88.2	88.1‡	84.1	83.9	84.9	83.1	81.4	82.1	81.9	81.0	82.0	+1.0
Taking amphetamines regularly <sup>d</sup>	95.4	94.2	95.6	94.9‡	92.9	93.9	93.2	93.0	92.2	92.2	92.0	92.8	93.8	+1.0
Trying sedatives (barbiturates) once or twice <sup>e</sup>	86.5	86.1	87.7	87.6	87.3	88.2	88.9	88.5	87.4	86.5	85.9	86.9	85.9	-1.0
Taking sedatives (barbiturates) regularly <sup>e</sup>	94.6	94.3	95.8	94.7	95.1	96.1	95.8	95.0	94.7	94.8	94.4	95.3	94.8	-0.4
Trying one or two drinks of an alcoholic beverage														
(beer, wine, liquor)	31.0	29.8	30.6	30.7	28.7	25.4	27.3	29.2	28.9	28.8	27.2	31.3	28.3	-3.0
Taking one or two drinks nearly every day	73.3	74.5	70.5	71.5	72.8	70.8	71.9	71.7	71.1	71.8	70.8	74.7	73.8	-0.9
Taking four or five drinks nearly every day	90.5	89.8	89.7	88.8	90.8	90.1	90.6	91.9	89.7	91.1	90.7	91.7	91.7	0.0
Having five or more drinks once or twice														
each weekend	68.8	68.9	67.6	68.8	70.0	70.1	71.6	72.6	71.9	74.2	72.5	75.8	72.5	-3.3
Smoking one or more packs of cigarettes per day	80.7	80.5	81.8	81.0	83.0	83.7	82.6	85.0	84.1	85.3	86.6	89.0	87.8	-1.3
Vape an e-liquid with nicotine ocasionally	_	_	_	_	_	_	_	_	_	_	62.0	59.2	58.8	-0.4
Vape an e-liquid with nicotine regularly <sup>f</sup>	_	_	_	_	_	_	_	_	_	_	71.8	70.9	70.4	-0.4
Use JUUL occasionally <sup>f</sup>	_	_	_	_	_	—	_	_	—	_	_	_	58.1	_
Use JUUL regularly <sup>f</sup>	_	_	_	_	_	_	_	_	_	_	_	_	69.2	_
Taking steroids	89.2	90.9	90.3	89.8	89.7	90.4	88.2	87.5	87.8	86.7	88.5	87.4	89.5	+2.1
Approximate weighted N =	2.450	2,314	2,233	2,449	2,384	2,301	2,147	2,078	2,193	2,000	1.870	1,918	1.946	

Table continued on next page.

## TABLE 14 (cont.)Trends in <u>Disapproval</u> of Drug Use in <u>Grade 12</u>

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. ' ‡ ' indicates that the question changed the following year. See relevant footnote for that drug. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

<sup>a</sup>The 1975 question asked about people who are 20 or older.

<sup>b</sup>Answer alternatives were: (1) Don't disapprove, (2) Disapprove, and (3) Strongly disapprove. Percentages are shown for categories (2) and (3) combined.

<sup>c</sup>Beginning in 2014 "molly" was added to the question on disapproval of using MDMA once or twice. 2014 and 2015 data are not comparable to earlier years due to this change. <sup>d</sup>In 2011 the list of examples was changed from upper, pep pill, bennie, speed to upper, speed, Adderall, Ritalin, etc. These changes likely explain the discontinuity in the 2011 results.

<sup>e</sup>In 2004 the question text was changed from barbiturates to sedatives/barbiturates and the list of examples was changed from downers, goofballs, reds, yellows, etc. to just downers. These changes likely explain the discontinuity in the 2004 results.

<sup>f</sup>Based on two of six forms; N is two times the N indicated.