

FIGURE 1
VAPING MARIJUANA
Trends in 30-Day Prevalence among College Students
and Youth not in College 1 to 4 Years beyond High School





FIGURE 2
VAPING NICOTINE
Trends in 30-Day Prevalence among College Students
and Youth not in College 1 to 4 Years beyond High School



Source. The Monitoring the Future study, the University of Michigan.

FIGURE 3
MARIJUANA

Trends in Annual Prevalence among College Students and Youth not in College 1 to 4 Years beyond High School





FIGURE 4
MARIJUANA

Trends in 30-Day Prevalence of Daily Use among College Students and Youth not in College 1 to 4 Years beyond High School



Source. The Monitoring the Future study, the University of Michigan.

FIGURE 5
ANY ILLICIT DRUG OTHER THAN MARIJUANA
Trends in Annual Prevalence among College Students
and Youth not in College 1 to 4 Years beyond High School





FIGURE 6
NARCOTICS OTHER THAN HEROIN^a
Trends in Annual Prevalence among College Students
and Youth not in College 1 to 4 Years beyond High School



Source. The Monitoring the Future study, the University of Michigan.

^aIn 2002 the question text was changed on three of the six questionnaire forms. The list of examples of narcotics other than heroin was updated: Talwin, laudanum, and paregoric—all of which had negligible rates of use by 2001—were replaced by Vicodin, OxyContin, and Percocet. The 2002 data presented here are based on the changed forms only. In 2003 the remaining forms were changed.

FIGURE 7
AMPHETAMINES^a
Trends in Annual Prevalence among College Students
and Youth not in College 1 to 4 Years beyond High School





FIGURE 8
MDMA (ECSTASY, MOLLY)^b
Trends in Annual Prevalence among College Students
and Youth not in College 1 to 4 Years beyond High School



Source. The Monitoring the Future study, the University of Michigan.

^aRevised questions about amphetamine use were introduced in 1982 to more completely exclude inappropriate reporting of nonprescription amphetamines. In 2013 the question wording was changed slightly in three of the six questionnaire forms. Data from 2013 are based on the changed forms only. The change was made to all forms in 2014.

^bIn 2014, a version of the question was added to one survey form that included "molly" in the description. In 2015, the remaining forms were changed to this updated wording.

FIGURE 9

LSD

Trends in Annual Prevalence among College Students and Youth not in College 1 to 4 Years beyond High School





FIGURE 10

COCAINE

Trends in Annual Prevalence among College Students and Youth not in College 1 to 4 Years beyond High School



Source. The Monitoring the Future study, the University of Michigan.

FIGURE 11

ALCOHOL

Trends in Two-Week Prevalence of Having 5 or More Drinks in a Row among College Students and Youth not in College 1 to 4 Years beyond High School





FIGURE 12

CIGARETTES

Trends in 30-Day Prevalence among College Students and Youth not in College 1 to 4 Years beyond High School



Source. The Monitoring the Future study, the University of Michigan.