

**TABLE 7**  
**Long-Term Trends in Thirty-Day Prevalence of Daily Use of Various Drugs for Twelfth Graders**

Percent who used daily in last thirty days

	Class of 1975	Class of 1976	Class of 1977	Class of 1978	Class of 1979	Class of 1980	Class of 1981	Class of 1982	Class of 1983	Class of 1984	Class of 1985	Class of 1986	Class of 1987	Class of 1988	Class of 1989	Class of 1990	Class of 1991	Class of 1992	Class of 1993	Class of 1994	Class of 1995	Class of 1996	Class of 1997	Class of 1998	Class of 1999	Class of 2000	'99-'00 change	
<i>Approx. N (in thousands) =</i>	9.4	15.4	17.1	17.8	15.5	15.9	17.5	17.7	16.3	15.9	16.0	15.2	16.3	16.3	16.7	15.2	15.0	15.8	16.3	15.4	15.4	14.3	15.4	15.2	13.6	12.8		
Marijuana/Hashish	6.0	8.2	9.1	10.7	10.3	9.1	7.0	6.3	5.5	5.0	4.9	4.0	3.3	2.7	2.9	2.2	2.0	1.9	2.4	3.6	4.6	4.9	5.8	5.6	6.0	6.0	0.0	
Inhalants <sup>c</sup>	—	*	*	0.1	*	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.1	0.2	0.2	0.3	0.2	0.1	0.1	0.1	0.1	0.2	0.1	0.2	0.2	0.2	0.0
<i>Inhalants, Adjusted<sup>e,d</sup></i>	—	—	—	—	0.1	0.2	0.2	0.2	0.2	0.2	0.4	0.4	0.4	0.3	0.3	0.3	0.5	0.2	0.2	—	—	0.4	0.2	0.9	0.3	0.3	-0.1	
<i>Amyl/Butyl Nitrites<sup>e,f</sup></i>	—	—	—	—	*	0.1	0.1	0.0	0.2	0.1	0.3	0.5	0.3	0.1	0.3	0.1	0.2	0.1	0.1	0.2	0.2	0.4	0.1	0.3	0.2	*	-0.2	
Hallucinogens	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.3	0.1	0.1	0.2	+0.1	
<i>Hallucinogens, Adjusted<sup>g</sup></i>	—	—	—	—	0.2	0.2	0.1	0.2	0.2	0.2	0.3	0.3	0.2	*	0.3	0.3	0.1	0.1	0.1	—	—	0.4	0.4	0.8	0.2	0.2	0.0	
LSD	*	*	*	*	*	*	0.1	*	0.1	0.1	0.1	*	0.1	*	*	0.1	0.1	0.1	0.1	0.1	0.1	*	0.2	0.1	0.1	0.1	+0.1	
<i>Hallucinogens Other Than LSD</i>	—	0.1	0.1	*	*	*	0.1	*	*	0.1	*	*	*	*	*	*	*	*	*	*	0.1	0.1	0.1	0.1	0.1	*	0.1	+0.1s
PCP <sup>e,f</sup>	—	—	—	—	0.1	0.1	0.1	0.1	0.1	0.1	0.3	0.2	0.3	0.1	0.2	0.1	0.1	0.1	0.1	0.3	0.3	0.3	0.1	0.3	0.2	0.2	0.0	
MDMA (Ecstasy) <sup>e</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.0	0.1	0.2	0.1	*	-0.1	
Cocaine	0.1	0.1	0.1	0.1	0.2	0.2	0.3	0.2	0.2	0.2	0.4	0.4	0.3	0.2	0.3	0.1	0.1	0.1	0.1	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.0	
Crack <sup>h</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.2	0.1	-0.1
Other Cocaine <sup>i</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	0.2	0.2	0.1	0.1	0.1	*	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.0
Heroin <sup>j</sup>	0.1	*	*	*	*	*	*	*	0.1	*	*	*	*	*	*	0.1	*	*	*	*	*	0.1	0.1	0.1	0.1	0.1	0.1	0.0
With a needle <sup>k</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.1	0.2	0.1	*	*	*	0.0	
Without a needle <sup>k</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	*	0.1	0.1	0.0	0.0	*	0.0	
Other Narcotics <sup>l</sup>	0.1	0.1	0.2	0.1	*	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.1	0.1	*	*	0.1	0.1	0.2	0.2	0.1	0.2	0.1	0.0	
Amphetamines <sup>b,l</sup>	0.5	0.4	0.5	0.5	0.6	0.7	1.2	0.7	0.8	0.6	0.4	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.3	0.3	0.3	0.5	+0.2s	
Methamphetamine <sup>m</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.1	0.1	0.0	
Crystal Meth. (Ice) <sup>m</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.1	0.1	0.1	0.1	*	0.1	0.1	0.1	*	*	0.1	0.0	
Sedatives <sup>l,n</sup>	0.3	0.2	0.2	0.2	0.1	0.2	0.2	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	0.1	0.1	0.1	0.1	0.1	0.2	0.1	-0.1s
Barbiturates <sup>l</sup>	0.1	0.1	0.2	0.1	*	0.1	0.1	0.1	0.1	*	0.1	0.1	0.1	*	0.1	0.1	0.1	*	0.1	*	0.1	0.1	0.1	0.1	0.1	0.2	0.1	-0.1s
Methaqualone <sup>l,n</sup>	*	*	*	*	*	0.1	0.1	0.1	*	*	*	*	*	0.1	*	*	*	*	0.1	0.0	0.1	0.1	0.0	0.1	0.0	0.0	0.0	0.0
Tranquilizers <sup>l</sup>	0.1	0.2	0.3	0.1	0.1	0.1	0.1	0.1	0.1	0.1	*	*	0.1	*	0.1	0.1	0.1	*	*	0.1	*	0.2	0.1	0.1	0.1	0.1	0.1	-0.1
Rohypno <sup>e</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.1	0.0	0.1	0.1	0.1	+0.1	
Alcohol																												
Daily <sup>o</sup>	5.7	5.6	6.1	5.7	6.9	6.0	6.0	5.7	5.5	4.8	5.0	4.8	4.8	4.2	4.2	3.7	3.6	3.4	2.5	—	—	—	—	—	—	—	—	
Been drunk daily <sup>m</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.9	0.8	0.9	1.2	1.3	1.6	2.0	1.5	1.9	1.7	-0.2	
5+ drinks in a row in last 2 weeks	36.8	37.1	39.4	40.3	41.2	41.4	40.5	40.8	38.7	36.7	36.8	37.5	34.7	33.0	32.2	29.8	27.9	27.5	28.2	29.8	30.2	31.3	31.5	30.8	30.0	-0.8		
Cigarettes																												
Daily	26.9	28.8	28.8	27.5	25.4	21.3	20.3	21.1	21.2	18.7	19.5	18.7	18.7	18.1	18.9	19.1	18.5	17.2	19.0	19.4	21.6	22.2	24.6	22.4	23.1	20.6	-2.5s	
Half-pack or more per day	17.9	19.2	19.4	18.8	16.5	14.3	13.5	14.2	13.8	12.3	12.5	11.4	11.4	10.6	11.2	11.3	10.7	10.0	10.9	11.2	12.4	13.0	14.3	12.6	13.2	11.3	-1.9ss	
Smokeless																												
Tobacco <sup>e,p</sup>	—	—	—	—	—	—	—	—	—	—	—	4.7	5.1	4.3	3.3	—	—	4.3	3.3	3.9	3.6	3.3	4.4	3.2	2.9	3.2	+0.3	
Steroids <sup>m</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.1	0.2	0.1	0.1	0.1	0.4	0.2	0.3	0.3	0.3	0.2	0.2	0.0	

NOTES: Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. \*\* indicates less than .05 percent but greater than 0 percent. See Table 4 for relevant footnotes. Any apparent inconsistency between the change estimate and the prevalence of use estimates for the two most recent classes is due to rounding error. Daily use is defined as use on 20 or more occasions in the past 30 days except for 5+ drinks, cigarettes, and smokeless tobacco, for which actual daily use is measured.

SOURCE: The Monitoring the Future Study, the University of Michigan.