

**TABLE 14**  
**Long-Term Trends in Lifetime Prevalence of Use of Various Drugs for Twelfth Graders**

	Percentage who ever used															
	Class of:															
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
Approx. N =	9,400	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200
Any Illicit Drug <sup>a,b</sup>	55.2	58.3	61.6	64.1	65.1	65.4	65.6	64.4	62.9	61.6	60.6	57.6	56.6	53.9	50.9	47.9
Any Illicit Drug Other Than Marijuana <sup>a,b,c</sup>	36.2	35.4	35.8	36.5	37.4	38.7	42.8	41.1	40.4	40.3	39.7	37.7	35.8	32.5	31.4	29.4
Marijuana/Hashish	47.3	52.8	56.4	59.2	60.4	60.3	59.5	58.7	57.0	54.9	54.2	50.9	50.2	47.2	43.7	40.7
Inhalants <sup>d</sup>	—	10.3	11.1	12.0	12.7	11.9	12.3	12.8	13.6	14.4	15.4	15.9	17.0	16.7	17.6	18.0
Inhalants, Adjusted <sup>d,e</sup>	—	—	—	—	18.2	17.3	17.2	17.7	18.2	18.0	18.1	20.1	18.6	17.5	18.6	18.5
Amyl/Butyl Nitrites <sup>f,g</sup>	—	—	—	—	11.1	11.1	10.1	9.8	8.4	8.1	7.9	8.6	4.7	3.2	3.3	2.1
Hallucinogens <sup>c</sup>	16.3	15.1	13.9	14.3	14.1	13.3	13.3	12.5	11.9	10.7	10.3	9.7	10.3	8.9	9.4	9.4
Hallucinogens, Adjusted <sup>c,h</sup>	—	—	—	—	17.7	15.6	15.3	14.3	13.6	12.3	12.1	11.9	10.6	9.2	9.9	9.7
LSD	11.3	11.0	9.8	9.7	9.5	9.3	9.8	9.6	8.9	8.0	7.5	7.2	8.4	7.7	8.3	8.7
Hallucinogens Other Than LSD <sup>c</sup>	14.1	12.1	11.2	11.6	10.7	9.8	9.1	8.0	7.3	6.6	6.5	5.7	5.4	4.1	4.3	4.1
PCP <sup>f,g</sup>	—	—	—	—	12.8	9.6	7.8	6.0	5.6	5.0	4.9	4.8	3.0	2.9	3.9	2.8
MDMA (Ecstasy) <sup>f</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Cocaine	9.0	9.7	10.8	12.9	15.4	15.7	16.5	16.0	16.2	16.1	17.3	16.9	15.2	12.1	10.3	9.4
Crack <sup>i</sup>	—	—	—	—	—	—	—	—	—	—	—	—	5.4	4.8	4.7	3.5
Other Cocaine <sup>j</sup>	—	—	—	—	—	—	—	—	—	—	—	—	14.0	12.1	8.5	8.6
Heroin <sup>k</sup>	2.2	1.8	1.8	1.6	1.1	1.1	1.1	1.2	1.2	1.3	1.2	1.1	1.2	1.1	1.3	1.3
With a needle <sup>l</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Without a needle <sup>l</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other Narcotics <sup>m,n</sup>	9.0	9.6	10.3	9.9	10.1	9.8	10.1	9.6	9.4	9.7	10.2	9.0	9.2	8.6	8.3	8.3
Amphetamines <sup>b,m</sup>	22.3	22.6	23.0	22.9	24.2	26.4	32.2†	27.9	26.9	27.9	26.2	23.4	21.6	19.8	19.1	17.5
Methamphetamine <sup>o</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Crystal Meth. (Ice) <sup>o</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2.7
Sedatives (Barbiturates) <sup>m</sup>	16.9	16.2	15.6	13.7	11.8	11.0	11.3	10.3	9.9	9.9	9.2	8.4	7.4	6.7	6.5	6.8
Sedatives, Adjusted <sup>m,p</sup>	18.2	17.7	17.4	16.0	14.6	14.9	16.0	15.2	14.4	13.3	11.8	10.4	8.7	7.8	7.4	7.5
Methaqualone <sup>m,q</sup>	8.1	7.8	8.5	7.9	8.3	9.5	10.6	10.7	10.1	8.3	6.7	5.2	4.0	3.3	2.7	2.3
Tranquillizers <sup>c,m</sup>	17.0	16.8	18.0	17.0	16.3	15.2	14.7	14.0	13.3	12.4	11.9	10.9	10.9	9.4	7.6	7.2
Rohypnol <sup>f</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Alcohol <sup>f</sup>	90.4	91.9	92.5	93.1	93.0	93.2	92.6	92.8	92.6	92.6	92.2	91.3	92.2	92.0	90.7	89.5
Been Drunk <sup>o</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Cigarettes	73.6	75.4	75.7	75.3	74.0	71.0	71.0	70.1	70.6	69.7	68.8	67.6	67.2	66.4	65.7	64.4
Smokeless Tobacco <sup>f,s</sup>	—	—	—	—	—	—	—	—	—	—	—	31.4	32.2	30.4	29.2	—
Steroids <sup>t</sup>	—	—	—	—	—	—	—	—	—	—	—	—	—	—	3.0	2.9

Cont'd

**TABLE 14 (cont'd)**  
**Long-Term Trends in Lifetime Prevalence of Use of Various Drugs for Twelfth Graders**

	Percentage who ever used																'05-'06 change
	Class of:																
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	
<i>Approx. N =</i>	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200	
Any Illicit Drug <sup>a,b</sup>	44.1	40.7	42.9	45.6	48.4	50.8	54.3	54.1	54.7	54.0	53.9	53.0	51.1	51.1	50.4	48.2	-2.2
Any Illicit Drug Other Than Marijuana <sup>a,b,c</sup>	26.9	25.1	26.7	27.6	28.1	28.5	30.0	29.4	29.4	29.0†	30.7	29.5	27.7	28.7	27.4	26.9	-0.4
Marijuana/Hashish	36.7	32.6	35.3	38.2	41.7	44.9	49.6	49.1	49.7	48.8	49.0	47.8	46.1	45.7	44.8	42.3	-2.5
Inhalants <sup>d</sup>	17.6	16.6	17.4	17.7	17.4	16.6	16.1	15.2	15.4	14.2	13.0	11.7	11.2	10.9	11.4	11.1	-0.2
Inhalants, Adjusted <sup>d,e</sup>	18.0	17.0	17.7	18.3	17.8	17.5	16.9	16.5	16.0	14.6	13.8	12.4	12.2	11.4	11.9	11.5	-0.4
Amyl/Butyl Nitrites <sup>f,g</sup>	1.6	1.5	1.4	1.7	1.5	1.8	2.0	2.7	1.7	0.8	1.9	1.5	1.6	1.3	1.1	1.2	0.0
Hallucinogens <sup>c</sup>	9.6	9.2	10.9	11.4	12.7	14.0	15.1	14.1	13.7	13.0†	14.7	12.0	10.6	9.7	8.8	8.3	-0.5
Hallucinogens, Adjusted <sup>c,h</sup>	10.0	9.4	11.3	11.7	13.1	14.5	15.4	14.4	14.2	13.6†	15.3	12.8	10.9	9.9	9.3	8.8	-0.5
LSD	8.8	8.6	10.3	10.5	11.7	12.6	13.6	12.6	12.2	11.1	10.9	8.4	5.9	4.6	3.5	3.3	-0.2
Hallucinogens Other Than LSD <sup>c</sup>	3.7	3.3	3.9	4.9	5.4	6.8	7.5	7.1	6.7	6.9†	10.4	9.2	9.0	8.7	8.1	7.8	-0.4
PCP <sup>f,g</sup>	2.9	2.4	2.9	2.8	2.7	4.0	3.9	3.9	3.4	3.4	3.5	3.1	2.5	1.6	2.4	2.2	-0.2
MDMA (Ecstasy) <sup>f</sup>	—	—	—	—	—	6.1	6.9	5.8	8.0	11.0	11.7	10.5	8.3	7.5	5.4	6.5	+1.1
Cocaine	7.8	6.1	6.1	5.9	6.0	7.1	8.7	9.3	9.8	8.6	8.2	7.8	7.7	8.1	8.0	8.5	+0.5
Crack <sup>l</sup>	3.1	2.6	2.6	3.0	3.0	3.3	3.9	4.4	4.6	3.9	3.7	3.8	3.6	3.9	3.5	3.5	+0.1
Other Cocaine <sup>l</sup>	7.0	5.3	5.4	5.2	5.1	6.4	8.2	8.4	8.8	7.7	7.4	7.0	6.7	7.3	7.1	7.9	+0.8
Heroin <sup>k</sup>	0.9	1.2	1.1	1.2	1.6	1.8	2.1	2.0	2.0	2.4	1.8	1.7	1.5	1.5	1.5	1.4	-0.1
With a needle <sup>l</sup>	—	—	—	—	0.7	0.8	0.9	0.8	0.9	0.8	0.7	0.8	0.7	0.7	0.9	0.8	-0.1
Without a needle <sup>l</sup>	—	—	—	—	1.4	1.7	2.1	1.6	1.8	2.4	1.5	1.6	1.8	1.4	1.3	1.1	-0.2
Other Narcotics <sup>m,n</sup>	6.6	6.1	6.4	6.6	7.2	8.2	9.7	9.8	10.2	10.6	9.9†	13.5	13.2	13.5	12.8	13.4	+0.6
Amphetamines <sup>b,m</sup>	15.4	13.9	15.1	15.7	15.3	15.3	16.5	16.4	16.3	15.6	16.2	16.8	14.4	15.0	13.1	12.4	-0.6
Methamphetamine <sup>o</sup>	—	—	—	—	—	—	—	—	8.2	7.9	6.9	6.7	6.2	6.2	4.5	4.4	-0.1
Crystal Meth. (Ice) <sup>o</sup>	3.3	2.9	3.1	3.4	3.9	4.4	4.4	5.3	4.8	4.0	4.1	4.7	3.9	4.0	4.0	3.4	-0.6
Sedatives (Barbiturates) <sup>m</sup>	6.2	5.5	6.3	7.0	7.4	7.6	8.1	8.7	8.9	9.2	8.7	9.5	8.8	9.9	10.5	10.2	-0.3
Sedatives, Adjusted <sup>m,p</sup>	6.7	6.1	6.4	7.3	7.6	8.2	8.7	9.2	9.5	9.3	8.9	10.2	9.1	10.1	11.0	10.6	-0.4
Methaqualone <sup>m,q</sup>	1.3	1.6	0.8	1.4	1.2	2.0	1.7	1.6	1.8	0.8	1.1	1.5	1.0	1.3	1.3	1.2	-0.1
Tranquilizers <sup>c,m</sup>	7.2	6.0	6.4	6.6	7.1	7.2	7.8	8.5	9.3	8.9†	10.3	11.4	10.2	10.6	9.9	10.3	+0.4
Rohypnol <sup>f</sup>	—	—	—	—	—	1.2	1.8	3.0	2.0	1.5	1.7	—	—	—	—	—	—
Alcohol <sup>f</sup>	88.0	87.5‡	80.0	80.4	80.7	79.2	81.7	81.4	80.0	80.3	79.7	78.4	76.6	76.8	75.1	72.7	-2.4 s
Been Drunk <sup>o</sup>	65.4	63.4	62.5	62.9	63.2	61.8	64.2	62.4	62.3	62.3	63.9	61.6	58.1	60.3	57.5	56.4	-1.0
Cigarettes	63.1	61.8	61.9	62.0	64.2	63.5	65.4	65.3	64.6	62.5	61.0	57.2	53.7	52.8	50.0	47.1	-2.9 s
Smokeless Tobacco <sup>f,s</sup>	—	32.4	31.0	30.7	30.9	29.8	25.3	26.2	23.4	23.1	19.7	18.3	17.0	16.7	17.5	15.2	-2.3
Steroids <sup>t</sup>	2.1	2.1	2.0	2.4	2.3	1.9	2.4	2.7	2.9	2.5	3.7	4.0	3.5	3.4	2.6	2.7	+0.1

Source: The Monitoring the Future Study, the University of Michigan.