

TABLE 4
Trends in Prevalence of Use of Smokeless Tobacco in Grades 8, 10, and 12

Cont.

	<u>1975</u>	<u>1976</u>	<u>1977</u>	<u>1978</u>	<u>1979</u>	<u>1980</u>	<u>1981</u>	<u>1982</u>	<u>1983</u>	<u>1984</u>	<u>1985</u>	<u>1986</u>	<u>1987</u>	<u>1988</u>	<u>1989</u>	<u>1990</u>	<u>1991</u>
Lifetime																	
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	22.2
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	28.2
12th Grade	—	—	—	—	—	—	—	—	—	—	—	31.4	32.2	30.4	29.2	—	—
30-Day																	
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	6.9
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	10.0
12th Grade	—	—	—	—	—	—	—	—	—	—	—	11.5	11.3	10.3	8.4	—	—
Daily																	
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1.6
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	3.3
12th Grade	—	—	—	—	—	—	—	—	—	—	—	4.7	5.1	4.3	3.3	—	—
Approximate Ns:																	
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	17,500
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	14,800
12th Grade	—	—	—	—	—	—	—	—	—	—	—	15,200	16,300	16,300	16,700	—	15,000

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: $s = .05$, $ss = .01$, $sss = .001$.

“—” indicates data not available.

Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

For 8th and 10th graders: Data based on one of two forms for 1991–1996 and on two of four forms beginning in 1997; N is one half of N indicated.

For 12th graders: Data based on one form; N is one fifth of N indicated in 1986–1988 and one sixth of N indicated beginning in 1989.

The prevalence of use of smokeless tobacco was not asked of 12th graders in 1990 and 1991. Prior to 1990 the prevalence-of-use question on smokeless tobacco was located near the end of one 12th-grade questionnaire form, whereas after 1991 the question was placed earlier and in a different form. This shift could explain the discontinuities between the corresponding data.

(Table continued on next page.)

TABLE 4 (cont.)
Trends in Prevalence of Use of Smokeless Tobacco in Grades 8, 10, and 12

	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>	<u>2006</u>	<u>2007</u>	<u>2008</u>	2007– 2008 change
Lifetime																		
8th Grade	20.7	18.7	19.9	20.0	20.4	16.8	15.0	14.4	12.8	11.7	11.2	11.3	11.0	10.1	10.2	9.1	9.8	+0.8
10th Grade	26.6	28.1	29.2	27.6	27.4	26.3	22.7	20.4	19.1	19.5	16.9	14.6	13.8	14.5	15.0	15.1	12.2	-2.9 ss
12th Grade	32.4	31.0	30.7	30.9	29.8	25.3	26.2	23.4	23.1	19.7	18.3	17.0	16.7	17.5	15.2	15.1	15.6	+0.5
30-Day																		
8th Grade	7.0	6.6	7.7	7.1	7.1	5.5	4.8	4.5	4.2	4.0	3.3	4.1	4.1	3.3	3.7	3.2	3.5	+0.3
10th Grade	9.6	10.4	10.5	9.7	8.6	8.9	7.5	6.5	6.1	6.9	6.1	5.3	4.9	5.6	5.7	6.1	5.0	-1.1
12th Grade	11.4	10.7	11.1	12.2	9.8	9.7	8.8	8.4	7.6	7.8	6.5	6.7	6.7	7.6	6.1	6.6	6.5	-0.1
Daily																		
8th Grade	1.8	1.5	1.9	1.2	1.5	1.0	1.0	0.9	0.9	1.2	0.8	0.8	1.0	0.7	0.7	0.8	0.8	-0.1
10th Grade	3.0	3.3	3.0	2.7	2.2	2.2	2.2	1.5	1.9	2.2	1.7	1.8	1.6	1.9	1.7	1.6	1.4	-0.2
12th Grade	4.3	3.3	3.9	3.6	3.3	4.4	3.2	2.9	3.2	2.8	2.0	2.2	2.8	2.5	2.2	2.8	2.7	-0.1
Approximate Ns:																		
8th Grade	18,600	18,300	17,300	17,500	17,800	18,600	18,100	16,700	16,700	16,200	15,100	16,500	17,000	16,800	16,500	16,100	15,700	
10th Grade	14,800	15,300	15,800	17,000	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800	16,400	16,200	16,200	16,100	15,100	
12th Grade	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200	14,500	14,000	

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001.

"—" indicates data not available.

Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

For 8th and 10th graders: Data based on one of two forms for 1991–1996 and on two of four forms beginning in 1997; *N* is one half of *N* indicated.

For 12th graders: Data based on one form; *N* is one fifth of *N* indicated in 1986–1988 and one sixth of *N* indicated beginning in 1989.

The prevalence of use of smokeless tobacco was not asked of 12th graders in 1990 and 1991. Prior to 1990 the prevalence-of-use question on smokeless tobacco was located near the end of one 12th-grade questionnaire form, whereas after 1991 the question was placed earlier and in a different form. This shift could explain the discontinuities between the corresponding data.