

TABLE 17
Long-Term Trends in 30-Day Prevalence of Use of Various Drugs in Grade 12

	Percentage who used in last 30 days																			
	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991			
Approximate weighted N =	9,400	15,400	17,100	17,800	15,500	15,900	17,500	17,700	16,300	15,900	16,000	15,200	16,300	16,300	16,700	15,200	15,000			
Any Illicit Drug ^{a,b}	30.7	34.2	37.6	38.9	38.9	37.2	36.9	32.5	30.5	29.2	29.7	27.1	24.7	21.3	19.7	17.2	16.4			
Any Illicit Drug other than Marijuana ^{a,b,c}	15.4	13.9	15.2	15.1	16.8	18.4	21.7	17.0	15.4	15.1	14.9	13.2	11.6	10.0	9.1	8.0	7.1			
Marijuana/Hashish	27.1	32.2	35.4	37.1	36.5	33.7	31.6	28.5	27.0	25.2	25.7	23.4	21.0	18.0	16.7	14.0	13.8			
Inhalants ^d	—	0.9	1.3	1.5	1.7	1.4	1.5	1.5	1.7	1.9	2.2	2.5	2.8	2.6	2.3	2.7	2.4			
Inhalants, Adjusted ^{d,e}	—	—	—	—	3.2	2.7	2.5	2.5	2.5	2.6	3.0	3.2	3.5	3.0	2.7	2.9	2.6			
Amyl/Butyl Nitrates ^{f,g}	—	—	—	—	2.4	1.8	1.4	1.1	1.4	1.4	1.6	1.3	1.3	0.6	0.6	0.6	0.4			
Hallucinogens ^c	4.7	3.4	4.1	3.9	4.0	3.7	3.7	3.4	2.8	2.6	2.5	2.5	2.5	2.2	2.2	2.2	2.2			
Hallucinogens, Adjusted ^{c,h}	—	—	—	—	5.3	4.4	4.5	4.1	3.5	3.2	3.8	3.5	2.8	2.3	2.9	2.3	2.4			
LSD	2.3	1.9	2.1	2.1	2.4	2.3	2.5	2.4	1.9	1.5	1.6	1.7	1.8	1.8	1.8	1.9	1.9			
Hallucinogens other than LSD ^c	3.7	2.3	3.0	2.7	2.4	2.3	2.1	1.7	1.5	1.6	1.3	1.3	1.1	0.7	0.8	0.8	0.7			
PCP ^{f,g}	—	—	—	—	2.4	1.4	1.4	1.0	1.3	1.0	1.6	1.3	0.6	0.3	1.4	0.4	0.5			
Ecstasy (MDMA) ^f	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—			
Cocaine	1.9	2.0	2.9	3.9	5.7	5.2	5.8	5.0	4.9	5.8	6.7	6.2	4.3	3.4	2.8	1.9	1.4			
Crack ⁱ	—	—	—	—	—	—	—	—	—	—	—	—	1.3	1.6	1.4	0.7	0.7			
Other Cocaine ^j	—	—	—	—	—	—	—	—	—	—	—	—	4.1	3.2	1.9	1.7	1.2			
Heroin ^k	0.4	0.2	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.3	0.3	0.2	0.2	0.2	0.3	0.2	0.2			
With a needle ^l	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—			
Without a needle ^l	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—			
Narcotics other than Heroin ^{m,n}	2.1	2.0	2.8	2.1	2.4	2.4	2.1	1.8	1.8	1.8	2.3	2.0	1.8	1.6	1.6	1.5	1.1			
Amphetamines ^{b,m}	8.5	7.7	8.8	8.7	9.9	12.1	15.8 [‡]	10.7	8.9	8.3	6.8	5.5	5.2	4.6	4.2	3.7	3.2			
Methamphetamine ^o	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—			
Crystal Methamphetamine (Ice) ^o	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.6	0.6		
Sedatives (Barbiturates) ^m	4.7	3.9	4.3	3.2	3.2	2.9	2.6	2.0	2.1	1.7	2.0	1.8	1.4	1.2	1.4	1.3	1.4			
Sedatives, Adjusted ^{m,p}	5.4	4.5	5.1	4.2	4.4	4.8	4.6	3.4	3.0	2.3	2.4	2.2	1.7	1.4	1.6	1.4	1.5			
Methaqualone ^{m,q}	2.1	1.6	2.3	1.9	2.3	3.3	3.1	2.4	1.8	1.1	1.0	0.8	0.6	0.5	0.6	0.2	0.2			
Tranquilizers ^{c,m}	4.1	4.0	4.6	3.4	3.7	3.1	2.7	2.4	2.5	2.1	2.1	2.1	2.0	1.5	1.3	1.2	1.4			
Rohypnol ^f	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—			
Alcohol ^f	68.2	68.3	71.2	72.1	71.8	72.0	70.7	69.7	69.4	67.2	65.9	65.3	66.4	63.9	60.0	57.1	54.0			
Been Drunk ^o	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	31.6			
Cigarettes	36.7	38.8	38.4	36.7	34.4	30.5	29.4	30.0	30.3	29.3	30.1	29.6	29.4	28.7	28.6	29.4	28.3			
Smokeless Tobacco ^{f,s}	—	—	—	—	—	—	—	—	—	—	—	11.5	11.3	10.3	8.4	—	—			
Steroids ^{m,t}	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.8	1.0	0.8			

Source. The Monitoring the Future study, the University of Michigan.

(Table continued on next page.)

TABLE 17 (cont.)
Long-Term Trends in 30-Day Prevalence of Use of Various Drugs for Grade 12

	Percentage who used in last 30 days																			2008– 2009 <u>change</u>
	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009		
Approximate weighted N =	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600	14,600	14,700	14,200	14,500	14,000	13,700		
Any Illicit Drug ^{a,b}	14.4	18.3	21.9	23.8	24.6	26.2	25.6	25.9	24.9	25.7	25.4	24.1	23.4	23.1	21.5	21.9	22.3	23.3	+0.9	
Any Illicit Drug other than Marijuana ^{a,b,c}	6.3	7.9	8.8	10.0	9.5	10.7	10.7	10.4	10.4‡	11.0	11.3	10.4	10.8	10.3	9.8	9.5	9.3	8.6	-0.6	
Marijuana/Hashish	11.9	15.5	19.0	21.2	21.9	23.7	22.8	23.1	21.6	22.4	21.5	21.2	19.9	19.8	18.3	18.8	19.4	20.6	+1.2	
Inhalants ^d	2.3	2.5	2.7	3.2	2.5	2.5	2.3	2.0	2.2	1.7	1.5	1.5	1.5	2.0	1.5	1.2	1.4	1.2	-0.2	
Inhalants, Adjusted ^{d,e}	2.5	2.8	2.9	3.5	2.9	2.9	3.1	2.4	2.4	2.1	1.8	2.3	1.9	2.3	1.7	1.6	1.5	1.8	+0.3	
Amyl/Butyl Nitrates ^{f,g}	0.3	0.6	0.4	0.4	0.7	0.7	1.0	0.4	0.3	0.5	0.6	0.7	0.7	0.5	0.3	0.5	0.3	0.6	+0.3	
Hallucinogens ^c	2.1	2.7	3.1	4.4	3.5	3.9	3.8	3.5	2.6‡	3.3	2.3	1.8	1.9	1.9	1.5	1.7	2.2	1.6	-0.6 s	
Hallucinogens, Adjusted ^{c,h}	2.3	3.3	3.2	4.6	3.8	4.1	4.1	3.9	3.0‡	3.5	2.7	2.7	2.2	2.5	1.8	2.1	2.6	1.9	-0.7 s	
LSD	2.0	2.4	2.6	4.0	2.5	3.1	3.2	2.7	1.6	2.3	0.7	0.6	0.7	0.7	0.6	0.6	1.1	0.5	-0.6 sss	
Hallucinogens other than LSD ^c	0.5	0.8	1.2	1.3	1.6	1.7	1.6	1.6	1.7‡	1.9	2.0	1.5	1.7	1.6	1.3	1.4	1.6	1.4	-0.3	
PCP ^{f,g}	0.6	1.0	0.7	0.6	1.3	0.7	1.0	0.8	0.9	0.5	0.4	0.6	0.4	0.7	0.4	0.5	0.6	0.5	0.0	
Ecstasy (MDMA) ^f	—	—	—	—	2.0	1.6	1.5	2.5	3.6	2.8	2.4	1.3	1.2	1.0	1.3	1.6	1.8	1.8	0.0	
Cocaine	1.3	1.3	1.5	1.8	2.0	2.3	2.4	2.6	2.1	2.1	2.3	2.1	2.3	2.3	2.5	2.0	1.9	1.3	-0.6 s	
Crack ⁱ	0.6	0.7	0.8	1.0	1.0	0.9	1.0	1.1	1.0	1.1	1.2	0.9	1.0	1.0	0.9	0.9	0.8	0.6	-0.2	
Other Cocaine ^j	1.0	1.2	1.3	1.3	1.6	2.0	2.0	2.5	1.7	1.8	1.9	1.8	2.2	2.0	2.4	1.7	1.7	1.1	-0.6 s	
Heroin ^k	0.3	0.2	0.3	0.6	0.5	0.5	0.5	0.5	0.7	0.4	0.5	0.4	0.5	0.5	0.4	0.4	0.4	0.4	0.0	
With a needle ^l	—	—	—	0.3	0.4	0.3	0.2	0.2	0.2	0.2	0.3	0.3	0.2	0.3	0.3	0.2	0.2	0.1	-0.1	
Without a needle ^l	—	—	—	0.6	0.4	0.6	0.4	0.4	0.7	0.3	0.5	0.4	0.3	0.5	0.3	0.4	0.2	0.3	+0.1	
Narcotics other than Heroin ^{m,n}	1.2	1.3	1.5	1.8	2.0	2.3	2.4	2.6	2.9	3.0‡	4.0	4.1	4.3	3.9	3.8	3.8	3.8	4.1	+0.2	
Amphetamines ^{b,m}	2.8	3.7	4.0	4.0	4.1	4.8	4.6	4.5	5.0	5.6	5.5	5.0	4.6	3.9	3.7	3.7	2.9	3.0	+0.2	
Methamphetamine ^o	—	—	—	—	—	—	—	1.7	1.9	1.5	1.7	1.7	1.4	0.9	0.9	0.6	0.6	0.5	0.0	
Crystal Methamphetamine (Ice) ^o	0.5	0.6	0.7	1.1	1.1	0.8	1.2	0.8	1.0	1.1	1.2	0.8	0.8	0.9	0.7	0.6	0.6	0.5	0.0	
Sedatives (Barbiturates) ^m	1.1	1.3	1.7	2.2	2.1	2.1	2.6	2.6	3.0	2.8	3.2	2.9	2.9	3.3	3.0	2.7	2.8	2.5	-0.3	
Sedatives, Adjusted ^{m,p}	1.2	1.3	1.8	2.3	2.3	2.1	2.8	2.8	3.1	3.0	3.4	3.0	2.9	3.5	3.1	2.8	2.9	2.6	-0.3	
Methaqualone ^{m,q}	0.4	0.1	0.4	0.4	0.6	0.3	0.6	0.4	0.2	0.5	0.3	0.4	0.5	0.5	0.4	0.4	0.2	0.3	0.0	
Tranquilizers ^{c,m}	1.0	1.2	1.4	1.8	2.0	1.8	2.4	2.5	2.6‡	2.9	3.3	2.8	3.1	2.9	2.7	2.6	2.7	0.0		
Rohypnol ^f	—	—	—	—	0.5	0.3	0.3	0.3	0.4	0.3	—	—	—	—	—	—	—	—		
Alcohol ^r	51.3‡	48.6	50.1	51.3	50.8	52.7	52.0	51.0	50.0	49.8	48.6	47.5	48.0	47.0	45.3	44.4	43.1	43.5	+0.3	
Been Drunk ^o	29.9	28.9	30.8	33.2	31.3	34.2	32.9	32.9	32.3	32.7	30.3	30.9	32.5	30.2	30.0	28.7	27.6	27.4	-0.2	
Cigarettes	27.8	29.9	31.2	33.5	34.0	36.5	35.1	34.6	31.4	29.5	26.7	24.4	25.0	23.2	21.6	21.6	20.4	20.1	-0.3	
Smokeless Tobacco ^{f,s}	11.4	10.7	11.1	12.2	9.8	9.7	8.8	8.4	7.6	7.8	6.5	6.7	6.7	7.6	6.1	6.6	6.5	8.4	+1.8	
Steroids ^{m,t}	0.6	0.7	0.9	0.7	0.7	1.0	1.1	0.9	0.8	1.3	1.4	1.3	1.6	0.9	1.1	1.0	1.0	1.0	0.0	

Source. The Monitoring the Future study, the University of Michigan.

Note. See relevant footnotes at the end of Table 15.