TABLE 6
Trends in <u>Disapproval</u> of Drug Use in <u>Grade 12</u>

Percentage who disapprove or strongly disapprove <sup>b</sup>

						Oroonia	9	попррі о		9-7						
Do you disapprove of people (who are 18 or older)																
loing each of the following? a	<u>1975</u>	1976	<u> 1977</u>	<u>1978</u>	1979	1980	<u>1981</u>	1982	1983	1984	1985	1986	1987	<u>1988</u>	1989	1990
se marijuana once or twice	47.0	38.4	33.4	33.4	34.2	39.0	40.0	45.5	46.3	49.3	51.4	54.6	56.6	60.8	64.6	67.8
se marijuana occasionally	54.8	47.8	44.3	43.5	45.3	49.7	52.6	59.1	60.7	63.5	65.8	69.0	71.6	74.0	77.2	80.5
e marijuana regularly	71.9	69.5	65.5	67.5	69.2	74.6	77.4	80.6	82.5	84.7	85.5	86.6	89.2	89.3	89.8	91.0
ring LSD once or twice	82.8	84.6	83.9	85.4	86.6	87.3	86.4	88.8	89.1	88.9	89.5	89.2	91.6	89.8	89.7	89.8
king LSD regularly	94.1	95.3	95.8	96.4	96.9	96.7	96.8	96.7	97.0	96.8	97.0	96.6	97.8	96.4	96.4	96.3
ing ecstasy (MDMA, Molly) once or twice c	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
ing cocaine once or twice	81.3	82.4	79.1	77.0	74.7	76.3	74.6	76.6	77.0	79.7	79.3	80.2	87.3	89.1	90.5	91.5
ring cocaine regularly	93.3	93.9	92.1	91.9	90.8	91.1	90.7	91.5	93.2	94.5	93.8	94.3	96.7	96.2	96.4	96.7
ring crack once or twice <sup>h</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	92.3
king crack occasionally <sup>h</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	94.3
king crack regularly <sup>h</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	94.9
ing cocaine powder once or twice <sup>h</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	87.9
king cocaine powder occasionally <sup>h</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	92.1
ing cocaine powder regularly <sup>h</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	93.7
ng heroin once or twice	91.5	92.6	92.5	92.0	93.4	93.5	93.5	94.6	94.3	94.0	94.0	93.3	96.2	95.0	95.4	95.1
ng heroin occasionally	94.8	96.0	96.0	96.4	96.8	96.7	97.2	96.9	96.9	97.1	96.8	96.6	97.9	96.9	97.2	96.7
ng heroin regularly	96.7	97.5	97.2	97.8	97.9	97.6	97.8	97.5	97.7	98.0	97.6	97.6	98.1	97.2	97.4	97.5
g heroin once or twice without using a needle	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
ng heroin occasionally without using a needle	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
g amphetamines once or twice <sup>d</sup>	74.8	75.1	74.2	74.8	75.1	75.4	71.1	72.6	72.3	72.8	74.9	76.5	80.7	82.5	83.3	85.3
ng amphetamines regularly <sup>d</sup>	92.1	92.8	92.5	93.5	94.4	93.0	91.7	92.0	92.6	93.6	93.3	93.5	95.4	94.2	94.2	95.5
g one or two drinks of an alcoholic beverage																
er, wine, liquor)	21.6	18.2	15.6	15.6	15.8	16.0	17.2	18.2	18.4	17.4	20.3	20.9	21.4	22.6	27.3	29.4
ng one or two drinks nearly every day	67.6	68.9	66.8	67.7	68.3	69.0	69.1	69.9	68.9	72.9	70.9	72.8	74.2	75.0	76.5	77.9
ng four or five drinks nearly every day	88.7	90.7	88.4	90.2	91.7	90.8	91.8	90.9	90.0	91.0	92.0	91.4	92.2	92.8	91.6	91.9
ng five or more drinks once or twice																
ch weekend	60.3	58.6	57.4	56.2	56.7	55.6	55.5	58.8	56.6	59.6	60.4	62.4	62.0	65.3	66.5	68.9
oking one or more packs of cigarettes per day	67.5	65.9	66.4	67.0	70.3	70.8	69.9	69.4	70.8	73.0	72.3	75.4	74.3	73.1	72.4	72.8
e marijuana occasionally <sup>e</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
e marijuana regularly <sup>e</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
e an e-liquid with nicotine occasionally <sup>e</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
e an e-liquid with nicotine regularly <sup>e</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_
king steroids	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	90.8
Approximate weighted N =	2,677	2,957	3,085	3,686	3,221	3,261	3,610	3,651	3,341	3,254	3,265	3,113	3,302	3,311	2,799	2,566

TABLE 6 (cont.)
Trends in <u>Disapproval</u> of Drug Use in <u>Grade 12</u>

Percentage who disapprove or strongly disapprove b

							Percenta	ge who c	lisapprov	e or stro	ongly dis	approve							i
Do you disapprove of people (who are 18 or older)																			
doing each of the following? <sup>a</sup>	<u>1991</u>	1992	1993	1994	<u>1995</u>	<u>1996</u>	<u>1997</u>	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	
Jse marijuana once or twice	68.7	69.9	63.3	57.6	56.7	52.5	51.0	51.6	48.8	52.5	49.1	51.6	53.4	52.7	55.0	55.6	58.6	55.5	
Jse marijuana occasionally	79.4	79.7	75.5	68.9	66.7	62.9	63.2	64.4	62.5	65.8	63.2	63.4	64.2	65.4	67.8	69.3	70.2	67.3	
Jse marijuana regularly	89.3	90.1	87.6	82.3	81.9	80.0	78.8	81.2	78.6	79.7	79.3	78.3	78.7	80.7	82.0	82.2	83.3	79.6	
Frying LSD once or twice	90.1	88.1	85.9	82.5	81.1	79.6	80.5	82.1	83.0	82.4	81.8	84.6	85.5	87.9	87.9	88.0	87.8	85.5	
aking LSD regularly	96.4	95.5	95.8	94.3	92.5	93.2	92.9	93.5	94.3	94.2	94.0	94.0	94.4	94.6	95.6	95.9	94.9	93.5	
rying ecstasy (MDMA, Molly) once or twice c	_	_	_	_	_	_	82.2	82.5	82.1	81.0	79.5	83.6	84.7	87.7	88.4	89.0	87.8	88.2	
rying cocaine once or twice	93.6	93.0	92.7	91.6	90.3	90.0	88.0	89.5	89.1	88.2	88.1	89.0	89.3	88.6	88.9	89.1	89.6	89.2	
aking cocaine regularly	97.3	96.9	97.5	96.6	96.1	95.6	96.0	95.6	94.9	95.5	94.9	95.0	95.8	95.4	96.0	96.1	96.2	94.8	
rying crack once or twice <sup>n</sup>	92.1	93.1	89.9	89.5	91.4	87.4	87.0	86.7	87.6	87.5	87.0	87.8	86.6	86.9	86.7	88.8	88.8	89.6	
aking crack occasionally h	94.2	95.0	92.8	92.8	94.0	91.2	91.3	90.9	92.3	91.9	91.6	91.5	90.8	92.1	91.9	92.9	92.4	93.3	
aking crack regularly <sup>n</sup>	95.0	95.5	93.4	93.1	94.1	93.0	92.3	91.9	93.2	92.8	92.2	92.4	91.2	93.1	92.1	93.8	93.6	93.5	
rying cocaine powder once or twice h	88.0	89.4	86.6	87.1	88.3	83.1	83.0	83.1	84.3	84.1	83.3	83.8	83.6	82.2	83.2	84.1	83.5	85.7	
aking cocaine powder occasionally h	93.0	93.4	91.2	91.0	92.7	89.7	89.3	88.7	90.0	90.3	89.8	90.2	88.9	90.0	89.4	90.4	90.6	91.7	
aking cocaine powder regularly h	94.4	94.3	93.0	92.5	93.8	92.9	91.5	91.1	92.3	92.6	92.5	92.2	90.7	92.6	92.0	93.2	92.6	92.8	
rying heroin once or twice	96.0	94.9	94.4	93.2	92.8	92.1	92.3	93.7	93.5	93.0	93.1	94.1	94.1	94.2	94.3	93.8	94.8	93.3	
aking heroin occasionally	97.3	96.8	97.0	96.2	95.7	95.0	95.4	96.1	95.7	96.0	95.4	95.6	95.9	96.4	96.3	96.2	96.8		Table conti
aking heroin regularly	97.8	97.2	97.5	97.1	96.4	96.3	96.4	96.6	96.4	96.6	96.2	96.2	97.1	97.1	96.7	96.9	97.1	95.9	on next pag
rying heroin once or twice without using a needle	_	_	_	_	92.9	90.8	92.3	93.0	92.6	94.0	91.7	93.1	92.2	93.1	93.2	93.7	93.6	94.2	
aking heroin occasionally without using a needle	_	_	_	_	94.7	93.2	94.4	94.3	93.8	95.2	93.5	94.4	93.5	94.4	95.0	94.5	94.9	95.3	
rying amphetamines once or twice <sup>d</sup>	86.5	86.9	84.2	81.3	82.2	79.9	81.3	82.5	81.9	82.1	82.3	83.8	85.8	84.1	86.1	86.3	87.3	87.2	
aking amphetamines regularly <sup>d</sup>	96.0	95.6	96.0	94.1	94.3	93.5	94.3	94.0	93.7	94.1	93.4	93.5	94.0	93.9	94.8	95.3	95.4	94.2	
rying one or two drinks of an alcoholic beverage																			
(beer, wine, liquor)	29.8	33.0	30.1	28.4	27.3	26.5	26.1	24.5	24.6	25.2	26.6	26.3	27.2	26.0	26.4	29.0	31.0	29.8	
aking one or two drinks nearly every day	76.5	75.9	77.8	73.1	73.3	70.8	70.0	69.4	67.2	70.0	69.2	69.1	68.9	69.5	70.8	72.8	73.3	74.5	
aking four or five drinks nearly every day	90.6	90.8	90.6	89.8	88.8	89.4	88.6	86.7	86.9	88.4	86.4	87.5	86.3	87.8	89.4	90.6	90.5	89.8	
Having five or more drinks once or twice																			
each weekend	67.4	70.7	70.1	65.1	66.7	64.7	65.0	63.8	62.7	65.2	62.9	64.7	64.2	65.7	66.5	68.5	68.8	68.9	
Smoking one or more packs of cigarettes per day	71.4	73.5	70.6	69.8	68.2	67.2	67.1	68.8	69.5	70.1	71.6	73.6	74.8	76.2	79.8	81.5	80.7	80.5	
/ape marijuana occasionally <sup>e</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
′ape marijuana regularly <sup>e</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
ape an e-liquid with nicotine occasionally <sup>e</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
ape an e-liquid with nicotine regularly <sup>e</sup>	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	_	
aking steroids	90.5	92.1	92.1	91.9	91.0	91.7	91.4	90.8	88.9	88.8	86.4	86.8	86.0	87.9	88.8	89.4	89.2	90.9	
Approximate weighted N =	2,547	2,645	2,723	2,588	2,603	2,399	2,601	2,545	2,310	2,150	2,144	2,160	2,442	2,455	2,460	2,377	2,450	2,314	

TABLE 6 (cont.)
Trends in <u>Disapproval</u> of Drug Use in <u>Grade 12</u>

Percentage who disapprove or strongly disapprove <sup>b</sup>

								- ''		3,								
Do you disapprove of people (who are 18 or older)																	2022–2023	
doing each of the following? <sup>a</sup>	2009	<u>2010</u>	<u>2011</u>	<u>2012</u>	<u>2013</u>	<u>2014</u>	<u>2015</u>	<u>2016</u>	<u>2017</u>	<u>2018</u>	2019pf	<u>2019e<sup>f</sup></u>	2020	2021 <sup>g</sup>	2022	2023	<u>change</u>	
Use marijuana once or twice	54.8	51.6	51.3	48.8	49.1	48.0	45.5	43.1	39.0	41.1	34.1	39.6	§	31.2*	35.0	34.6	-0.4	
Use marijuana occasionally	65.6	62.0	60.9	59.1	58.9	56.7	52.9	50.5	46.7	49.2	41.4	46.6	§	38.6*	41.6	39.8	-1.7	
Use marijuana regularly	80.3	77.7	77.5	77.8	74.5	73.4	70.7	68.5	64.7	66.7	63.4	66.7	§	58.0*	61.6	59.7	-1.9	
Trying LSD once or twice	88.2	86.5	86.3	87.2	86.6	85.0	81.7	82.4	78.0	80.5	76.1	77.7	§	68.7*	72.8	72.7	-0.1	
Taking LSD regularly	95.3	94.3	94.9	95.2	95.3	94.7	92.5	92.4	92.7	93.4	93.8	92.8	§	90.3*	89.8	90.7	+0.9	
Trying ecstasy (MDMA, Molly) once or twice <sup>c</sup>	88.2	86.3	83.9	87.1	84.9‡	83.1	84.5	84.0	85.1	85.6	89.8	87.6	§	85.5*	86.6	85.9	-0.7	
Trying cocaine once or twice	90.8	90.5	91.1	91.0	92.3	90.0	89.0	88.4	88.0	88.9	88.5	88.8	§	81.7*	88.7	87.0	-1.7	
Taking cocaine regularly	96.5	96.0	96.0	96.8	96.7	96.3	95.2	94.8	94.8	95.8	96.5	95.8	§	92.6*	95.0	95.1	0.0	
Trying crack once or twice <sup>h</sup>	90.9	89.8	91.4	92.8	91.4	89.3	90.2	90.1	89.7	90.4	88.7	85.1	§	87.4*	87.1	86.1	-1.0	
Taking crack occasionally <sup>h</sup>	94.0	92.6	93.9	95.0	93.6	91.9	92.5	92.0	91.8	92.2	91.1	85.7	§	90.1*	88.4	89.8	+1.4	
Taking crack regularly <sup>h</sup>	94.3	93.1	94.4	95.4	94.1	92.4	92.8	92.6	92.5	92.5	91.5	85.0	§	90.1*	88.9	90.3	+1.4	
Trying cocaine powder once or twice <sup>h</sup>	87.3	87.0	88.1	88.7	88.2	85.5	86.4	86.6	85.5	86.5	85.7	82.5	§	83.1*	83.4	84.3	+0.8	
Гaking cocaine powder occasionally <sup>h</sup>	92.3	91.0	92.2	93.0	91.7	90.4	91.3	90.6	90.3	91.3	90.1	84.3	§	86.6*	86.1	88.2	+2.1	
Гаking cocaine powder regularly <sup>h</sup>	93.9	92.6	93.8	95.0	94.1	91.7	92.4	92.0	92.2	92.0	91.2	85.6	§	89.5*	89.2	90.5	+1.3	
Trying heroin once or twice	94.7	93.9	94.3	95.8	95.6	94.7	94.2	94.1	93.7	95.0	95.7	93.9	§	92.8*	92.9	92.3	-0.6	
Taking heroin occasionally	96.9	96.2	96.3	97.0	96.9	96.6	95.3	95.5	95.5	96.4	96.7	95.9	§	94.9*	95.7	95.1	-0.6	Table continue
Taking heroin regularly	97.4	96.4	96.7	97.4	97.4	97.1	96.4	95.7	95.9	96.8	97.3	96.3	§	96.3*	96.7	95.6	-1.1	on next page.
Trying heroin once or twice without using a needle	94.7	93.2	92.6	95.2	93.7	92.5	92.6	93.8	93.3	93.0	95.2	95.0	§	93.4*	93.1	94.5	+1.3	
Faking heroin occasionally without using a needle	95.5	94.5	94.1	95.9	94.6	93.5	92.8	94.0	93.8	93.4	95.4	95.1	§	93.9*	93.8	94.7	+0.9	
Trying amphetamines once or twice <sup>d</sup>	88.2	88.1‡	84.1	83.9	84.9	83.1	81.4	82.1	81.9	81.0	80.3	83.5	§	78.5*	84.0	82.6	-1.3	
Taking amphetamines regularly <sup>d</sup>	95.6	94.9‡	92.9	93.9	93.2	93.0	92.2	92.2	92.0	92.8	94.4	93.3	§	88.3*	91.2	91.3	+0.1	
Trying one or two drinks of an alcoholic beverage																		
(beer, wine, liquor)	30.6	30.7	28.7	25.4	27.3	29.2	28.9	28.8	27.2	31.3	26.3	30.1	§	22.3*	26.7	26.5	-0.3	
Taking one or two drinks nearly every day	70.5	71.5	72.8	70.8	71.9	71.7	71.1	71.8	70.8	74.7	73.4	74.1	§	67.4*	71.0	72.4	+1.4	
Taking four or five drinks nearly every day	89.7	88.8	90.8	90.1	90.6	91.9	89.7	91.1	90.7	91.7	91.5	91.9	§	91.8*	92.2	92.9	+0.7	
Having five or more drinks once or twice																		
each weekend	67.6	68.8	70.0	70.1	71.6	72.6	71.9	74.2	72.5	75.8	75.0	70.2	§	57.8*	66.9	66.0	-0.8	
Smoking one or more packs of cigarettes per day	81.8	81.0	83.0	83.7	82.6	85.0	84.1	85.3	86.6	89.0	87.9	87.7	§	86.5*	86.3	87.4	+1.2	
/ape marijuana occasionally <sup>e</sup>	_	_	_	_	_	_	_	_	_	_	_	_	§	48.0*	52.8	57.5	+4.7 s	
Vape marijuana regularly <sup>e</sup>	_	_	_	_	_	_	_	_	_	_	_	_	§	64.5*	68.3	69.3	+0.9	
Vape an e-liquid with nicotine occasionally <sup>e</sup>	_	_	_	_	_	_	_	_	62.0	59.2	56.6	60.7	§	60.3*	64.9	69.3	+4.4 s	
Vape an e-liquid with nicotine regularly <sup>e</sup>	_	_	_	_	_	_	_	_	71.8	70.9	70.1	70.7	§	73.2*	76.0	79.6	+3.6	
Taking steroids	90.3	89.8	89.7	90.4	88.2	87.5	87.8	86.7	88.5	87.4	88.7	90.3	§	80.9*	84.5	81.9	-2.7	
Approximate weighted N =	2,233	2,449	2.384	2.301	2.147	2.078	2.193	2.000	1.870	1.918	876	975	8	1,441	1,539	1.220		

## TABLE 6 (cont.) Trends in Disapproval of Drug Use in Grade 12

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .001. '—' indicates data not available. '‡' indicates that the question

changed the following year. See relevant footnote for that drug. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

§Estimates not presented due to insufficient data this year.

\*Comparison of 2021+ estimates with previous years may be subject to a survey mode effect in 2019. The size and direction of the mode effect (if any) is indicated by the difference between the estimates in the '2019p' and the '2019e' columns. The '2019p' column reports estimates based on students in the randomly-selected half of schools that used paper-and-pencil questionnaires (used in 2018 and all previous years). The '2019e' column reports estimates on the other half that used electronic data collection on devices connected to the internet (used in 2021 and all subsequent years).

The 1975 question asked about people who are 20 or older.

bAnswer alternatives were: (1) Don't disapprove, (2) Disapprove, and (3) Strongly disapprove. Percentages are shown for categories (2) and (3) combined.

<sup>6</sup>Beginning in 2014 "molly" was added to the question on disapproval of using MDMA once or twice. 2014 and 2015 data are not comparable to earlier years due to this change.

<sup>d</sup>In 2011 the list of examples was changed from upper, pep pill, bennie, speed to upper, speed, Adderall, Ritalin, etc. These changes likely explain the discontinuity in the 2011 results.

<sup>e</sup>Based on two of six forms; N is two times the N indicated.

<sup>1</sup>The '2019p' column reports estimates from students in the randomly-selected half of schools that completed the 2019 questionnaire using paper and pencil. The '2019e' column reports estimates for the other half in schools that completed the 2019 questionnaire using web-connected electronic tablets. Estimates in italics indicate statistically significant (p<.05) differences in 2019 between the two survey modes.

<sup>9</sup>Sample is decreased by approximately 50% for the following drugs due to survey question experiments: amphetamines, cocaine, alcohol, vaping nicotine, vaping marijuana, heroin without using a needle, Ecstasy (MDMA, molly), and JUUL.

hThis estimate based on a question that was placed in a different form starting in 2021. Results from each form are nationally representative by themselves, as well as when combined.